

Estimating scale-specific and localized spatial patterns in allele frequency

Jesse R. Lasky^{1,2}, Margarita Takou¹, Diana Gamba¹, and Timothy H. Keitt³

¹Department of Biology, Pennsylvania State University, University Park, PA 16802, USA

²Correspondence: lasky@psu.edu

³Department of Integrative Biology, University of Texas at Austin, Austin, TX 78712, USA

May 4, 2024

Abstract

Characterizing spatial patterns in allele frequencies is fundamental to evolutionary biology because these patterns contain evidence of underlying processes. However, the spatial scales at which gene flow, changing selection, and drift act are often unknown. Many of these processes can operate inconsistently across space, causing non-stationary patterns. We present a wavelet approach to characterize spatial pattern in allele frequency that helps solve these problems. We show how our approach can characterize spatial patterns in relatedness at multiple spatial scales, i.e. a multi-locus wavelet genetic dissimilarity. We also develop wavelet tests of spatial differentiation in allele frequency and quantitative trait loci (QTL). With simulation we illustrate these methods under different scenarios. We also apply our approach to natural populations of *Arabidopsis thaliana* to characterize population structure and identify locally-adapted loci across scales. We find, for example, that *Arabidopsis* flowering time QTL show significantly elevated genetic differentiation at 300 to 1300 km scales. Wavelet transforms of allele frequencies offer a flexible way to reveal geographic patterns and underlying evolutionary processes.

Running head: Wavelet-transformed spatial genetic patterns

Keywords: Landscape genetics, F_{ST} , local adaptation, isolation by distance

1 Introduction

2 Geographic clines in allele frequency are a classic pattern in evolutionary bi-
3 ology, being frequently observed in nature and having extensive theory for the
4 underlying processes. For example, theory describes how limited gene flow and
5 drift (Wright 1931) or changing selection (Haldane 1948) can generate allele fre-
6 quency differences between populations. Accordingly, researchers often estimate
7 and model spatial allele frequency patterns to make inferences about underly-
8 ing evolutionary and ecological mechanisms. To do so, researchers often divide
9 sampled individuals into discrete groups (populations) among which differences
10 in allele frequencies are calculated. A common such approach involves estimat-
11 ing F_{ST} , the proportion of total allele frequency variation that differs between
12 discrete populations (Lewontin and Krakauer 1973; Wright 1949).

13 However, many species exist as more or less continuously distributed popu-
14 lations. Theoretical study of allele frequency change across continuous popu-
15 lations began as early as Wright (1943) and Malécot (1948), who found expec-
16 tations for genetic differentiation or kinship as functions of gene flow and geo-
17 graphic distance. Later progress included diffusion models (Nagylaki 1978) and
18 stepping stone/lattice models (Kimura and Weiss 1964) giving expectations for
19 correlation in allele frequencies across distance, and models accounting for pop-
20 ulation regulation by negative density dependence (Nick H. Barton, Depaulis,
21 and Etheridge 2002).

22 Despite these theoretical advances, the statistical tools for inference on con-
23 tinuously distributed populations have lagged (Bradburd and Peter L. Ralph
24 2019; Hancock, Toczydlowski, and Bradburd 2023). Nevertheless, statistical ap-
25 proaches to studying spatial pattern in continuous populations include models
26 relating landscape features to gene flow (McRae et al. 2008), calculating cor-
27 relations between spatial functions and genotype (Wagner, Chávez-Pesqueira,
28 and Forester 2017; Yang et al. 2012), and applying discrete landscape grids to
29 identify geographic regions where genetic turnover is particularly high or low
30 (Petkova, Novembre, and Stephens 2016). Approaches have been developed
31 to estimate the average distance of gene flow from the slope of genetic diver-
32 gence versus geographic distance (Rousset 2000; X. Vekemans and O. J. Hardy
33 2004), to estimate localized genetic "neighborhoods" (Shirk and Cushman 2014;
34 Wright 1946), and to model both discrete and continuous relatedness patterns
35 simultaneously (Bradburd, G. M. Coop, and Peter L. Ralph 2018).

36 In recent years researchers have collected many large, broadly distributed
37 DNA sequence datasets from diverse species (Alonso-Blanco et al. 2016; Machado
38 et al. 2021; J. Wang et al. 2020; Yeaman et al. 2016). Statistical inference can
39 be applied to these data to understand gene flow, demographic histories, and
40 spatially-varying selection. Despite the progress made by previous approaches,
41 there remain challenges.

42 1.1 The form and scale of relevant spatial patterns is un- 43 known

44 Humans can infer seemingly meaningful patterns in even randomly generated
45 images (Ayton and Fischer 2004; Blakemore et al. 2003; Fyfe et al. 2008). So
46 what are the spatial patterns we are looking for? The functional forms (i.e.
47 shapes) of both spatially-varying selection and neutral processes (e.g. dispersal
48 kernels) are often unknown, as are the forms of resulting spatial patterns. For
49 example, the specific environmental gradients driving changing selection are
50 often not known, nor is the spatial scale at which they act, and whether they
51 change at the same rate consistently across a landscape.

52 In the case of neutral processes, a homogeneous landscape approximately
53 at equilibrium is rarely of interest to empiricists. Instead, the influence of het-
54 erogeneous landscapes (Manel et al. 2003) and historical contingency is usually
55 a major force behind spatial patterns in allele frequency and traits (Excoffier
56 and Ray 2008). As a result, researchers often attempt to characterize spatial
57 patterns of relatedness and genetic similarity to make inferences about varia-
58 tion in gene flow (McRae et al. 2008; Peterman 2018; I. J. Wang, Savage, and
59 Bradley Shaffer 2009) and recent population expansion (Slatkin 1993). The in-
60 fluence of gene flow, drift, and range expansion can occur at a variety of spatial
61 scales, and in different ways across a heterogenous landscape. For example, the
62 rate at which relatedness decays over geographic distance can change abruptly
63 at major barriers (Rosenberg et al. 2005). However, the scale-specificity and
64 non-stationarity of such patterns can be challenging to characterize.

65 1.2 The spatially-varying selective gradients causing local 66 adaptation are unknown

67 One important force behind allele frequency clines is changing selection due
68 to environmental gradients, resulting in local adaptation. However, it is often
69 not clear what environmental gradients drive local adaptation (Kawecki and
70 Ebert 2004). This is especially true of non-model systems and those with little
71 existing natural history knowledge. Even for well-studied species, it is not trivial
72 to identify the specific environmental conditions that change in space and drive
73 local adaptation. Ecology is complex, and abiotic and biotic conditions are high-
74 dimensional. Rather than *a priori* selection of a putative selective gradient,
75 an alternative approach is to search for spatial patterns in allele frequencies
76 that cannot be explained by neutral processes. This approach is embodied
77 by several statistics and approaches, such as F_{ST} (Weir and Cockerham 1984),
78 XtX (Gautier 2015), spatial ancestry analysis (SPA) (Yang et al. 2012), Moran's
79 eigenvector maps (MEMs) (Wagner, Chávez-Pesqueira, and Forester 2017), and
80 others.

1.3 Many approaches rely on discretization of population boundaries

Some of the aforementioned approaches rely on dividing sampled individuals into discrete spatial groups. F_{ST} is one such approach, that was introduced by Wright (1949) and defined as the "correlation between random gametes, drawn from the same subpopulation, relative to the total", where the definition of "total" has been interpreted differently by different authors (Bhatia et al. 2013). The classic approach of calculating F_{ST} to test for selection was usually applied to a small number of locations, a situation when discretization (i.e. deciding which individuals genotyped belong in which population) was a simpler problem. Current studies often sample and sequence individuals from hundreds of locations, and so the best approach for discretizing these genotyped individuals into defined 'populations' is less clear. In addition to the spatial scale of subpopulations, at issue is precisely where to place the boundaries between populations. The problem is enhanced for broadly distributed species, connected by gene flow, that lack clear spatially distinct populations (Emily B. Josephs et al. 2019). Even if clustering algorithms appear to show clustering of genotypes, these methods can be sensitive to sampling bias (e.g. geographic clustering) and can mislead as to the existence of discrete subpopulations (Frantz et al. 2009; Serre and Pääbo 2004).

Some approaches are not limited by discretization, and might be generally termed "population-agnostic" because discrete populations are not defined. These instead use ordination of genetic loci or geographic location. Approaches that use ordination (such as PCA) of genetic loci look for particular loci with strong loadings on PCs (Duforet-Frebourg et al. 2016) or traits with an unexpectedly high correlation with individual PCs (Emily B. Josephs et al. 2019). Alternatively, ordination of distance or spatial neighborhood matrices can create spatial functions that can be used in correlation tests with genetic loci (Wagner, Chávez-Pesqueira, and Forester 2017). However, ordinations to create individual rotated axes are not done with respect to biology and so might not be ideal for characterizing biological patterns. For example, ordinations of genetic loci are heavily influenced by global outliers of genetic divergence (Peter, Petkova, and Novembre 2020) and uneven sampling (McVean 2009). Ordinations like PCA also often lack parametric null distributions for hypothesis testing.

1.4 Wavelet characterization of spatial pattern

Instead of discretizing sampled locations into populations, one could model allele frequencies with flexible but smooth functions. Wavelet transforms allow one to characterize the location and the scale or frequency of a signal (Daubechies 1992). Daubechies (1992) gives a nice analogy of wavelet transforms: they are akin to written music, which indicates a signal of a particular frequency (musical notes of different pitch) at a particular location (the time at which the note is played, in the case of music). Applying this analogy to genetics, the frequency is the rate at which allele frequencies change in space, and the location

124 is the part of a landscape where allele frequencies change at this rate. Applying
 125 wavelet basis functions to spatial genetic data could allow us to characterize
 126 localized patterns in allele frequency, and dilating the scale of these functions
 127 could allow us to characterize scale-specific patterns in allele frequency (see
 128 Figure S1 in File S1 for an example). Note that wavelets are distinct from
 129 Fourier analysis. Wavelets capture localized signals because the basis functions'
 130 variance goes to zero moving away from the focal location, while Fourier can
 131 only capture global average patterns as it uses stationary (unchanging) basis
 132 functions. Wavelet transforms have had some recent applications in modeling
 133 ancestry along the genome (Groh and G. Coop 2023; Pugach et al. 2011) but
 134 have not been implemented to model geographic genetic patterns.

135 Keitt (2007) created a wavelet approach for characterizing spatial patterns
 136 in ecological communities. He used this approach to identify locations and scales
 137 with particular high community turnover, and created null-hypothesis testing of
 138 these patterns. These spatial patterns in the abundance of multiple species are
 139 closely analogous to spatial patterns in allele frequency of many genetic markers
 140 across the genome, and previous spatial genetic studies have also profited by
 141 borrowing tools from spatial community ecology (Fitzpatrick and Keller 2015;
 142 Jesse R. Lasky, Des Marais, et al. 2012). Here we modify and build on this
 143 approach to characterize spatial pattern in allele frequency across the genome
 144 and at individual loci.

145 2 Methods

146 2.1 Wavelet characterization of spatial pattern in allele 147 frequency

148 Our implementation here begins by following the work of Keitt (2007) in char-
 149 acterizing spatial community turnover, except that we characterize genomic
 150 patterns using allele frequencies of multiple loci in place of abundances of mul-
 151 tiple species in ecological communities. In later sections of this paper we build
 152 off this approach and develop new tests for selection on specific loci. Our im-
 153 plementation of wavelets allows estimation of scale-specific signals (here, allele
 154 frequency clines) centered on a given point, a, b , in two-dimensional space. We
 155 use a version of the Difference-of-Gaussians (DoG) wavelet function (Figure S1
 156 in File S1) (Muraki 1995). We start with a Gaussian smoothing function cen-
 157 tered at a, b for a set of sampling points $\Omega = \{(u_1, v_1), (u_2, v_2), \dots, (u_n, v_n)\}$,
 158 which takes the form

$$\eta_{a,b}^s(x, y) = \frac{k\left(\frac{x-a}{s}, \frac{y-b}{s}\right)}{\sum_{(u,v) \in \Omega} k\left(\frac{u-a}{s}, \frac{v-b}{s}\right)}, \quad (1)$$

159 where s controls the scale of analysis and $k(x, y)$ is the Gaussian kernel
 160 $k(x, y) = e^{-(x^2+y^2)/2}$.

161 The DoG wavelet function then takes the form

$$\eta_{a,b}^s(x, y) = \eta_{a,b}^s(x, y) - \eta_{a,b}^{\beta s}(x, y) \quad (2)$$

162 where $\beta > 1$, and so the larger scale smooth function is subtracted from
 163 the smaller scale smooth to characterize the scale-specific pattern. If we use
 164 $\beta = 1.87$, then the dominant scale of analysis resulting from the DoG is s
 165 distance units (Keitt 2007). This formulation of the wavelet kernel is similar in
 166 shape to the derivative-of-Gaussian kernel and has the advantage of maintaining
 167 admissibility (Daubechies 1992) even near boundaries, as each of the smoothing
 168 kernels $\eta_{a,b}^s$ are normalized over the samples such that their difference integrates
 169 to zero.

170 Let $f_i(u, v)$ be the major allele frequency of the i th locus from a set of
 171 I biallelic markers at a location with spatial coordinates u, v . The adaptive
 172 wavelet transform of allele frequency data at locus i , centered at a, b and at
 173 scale s is then

$$(T^{wav} f_i)(a, b, s) = \frac{1}{h_{a,b}(s)} \sum_{(u,v) \in \Omega} \psi_{a,b}^s(u, v) f_i(u, v), \quad (3)$$

174 where the right summation is of the product of the smooth function and
 175 the allele frequencies across locations. The magnitude of this summation will
 176 be greatest when the DoG wavelet filter matches the allele frequency cline.
 177 That is, when the shape of the wavelet filter matches the allele frequency cline
 178 in space, the product of $\psi_{a,b}^s(u, v)$ and $f_i(u, v)$ will resonate (increase in ampli-
 179 tude) yielding greater variation among locations in $(T^{wav} f_i)(a, b, s)$, the wavelet-
 180 transformed allele frequencies. When the spatial pattern in the wavelet filter
 181 and allele frequencies are discordant, the variation in their product, and hence
 182 the wavelet-transformed allele frequency, is reduced. For consistency, here we
 183 choose major allele frequency for $f_i(u, v)$, though in practice the signing of alleles
 184 has little impact on our results.

185 The $h_{a,b}(s)$ term in equation 3 is used to normalize the variation in the
 186 wavelet function so that the wavelet transforms $T^{wav} f_i$ are comparable for dif-
 187 ferent scales s and locations a, b :

$$h_{a,b}(s) = \sqrt{\sum_{(u,v) \in \Omega} [\psi_{a,b}^s(u, v)]^2} \quad (4)$$

188 . When a, b is far from locations in Ω relative to the scale s , the Gaussian
 189 functions $[\eta_{a,b}^s(x, y)]$ that make up the wavelet function ψ are only evaluated
 190 over a range where they remain close to zero. Thus unsampled geographic
 191 regions will have very small $h_{a,b}(s)$, the term used to normalize for local variation
 192 in the wavelet basis functions. In turn, very small $h_{a,b}(s)$ dramatically and
 193 undesirably inflates the wavelet transformed allele frequencies (equation 3) in
 194 these geographic regions where there is little sampling relative to s . For this
 195 reason we do not calculate the wavelet transform for locations a, b where there
 196 are no locations sampled closer than $2s$ distance units.

197 Below we illustrate how to apply this wavelet transform (equation 3) of
 198 spatial allele frequency patterns to characterize genome-wide patterns, as well
 199 as to test for local adaption at individual loci.

200 **2.1.1 Wavelet characterization of spatial pattern in multiple loci**

201 Researchers are often interested in characterizing spatial patterns aggregated
 202 across multiple loci across the genome to understand patterns of relatedness,
 203 population structure, and demographic history. Here, we specifically want to
 204 characterize heterogeneity in spatial patterns, because this heterogeneity in pat-
 205 tern may reflect heterogeneity in underlying processes: where there is hetero-
 206 geneity in migration rates, such as where there are migration barriers (Petkova,
 207 Novembre, and Stephens 2016), or where there are recent range expansions such
 208 that spatial patterns are farther from equilibrium (Slatkin 1993).

209 We use

$$D_{a,b}^{wav}(s) = \sqrt{\sum_{i=1}^I [(T^{wav} f_i)(a, b, s)]^2} \quad (5)$$

210 to calculate a "wavelet genetic distance" or "wavelet genetic dissimilarity."
 211 This wavelet genetic dissimilarity is computed as the euclidean distance (in the
 212 space of allele frequencies across the genome) between the genetic composition
 213 centered at a, b and other locations across s distance units. This wavelet genetic
 214 dissimilarity $D_{a,b}^{wav}(s)$ is localized in space and scale-specific. This quantity cap-
 215 tures the level of genetic turnover at scale s centered at a, b , and is capturing
 216 similar information as the increase in average genetic distance between a geno-
 217 type at a, b and other genotypes s distance units away. To obtain the average
 218 dissimilarity across the landscape, one can also calculate the mean of $D_{a,b}^{wav}(s)$
 219 across locations a, b at each sampled site, to get a mean wavelet genetic dissim-
 220 ilarity for s . A benefit of using the wavelet transformation over sliding window
 221 approaches (e.g. Bishop, Chambers, and I. J. Wang 2023) is that wavelets
 222 smoothly incorporate patterns from samples that are not precisely s distance
 223 units away and can be centered at any location of the analyst's choosing.

224 **2.1.2 Testing the null hypothesis of no spatial pattern in allele fre-** 225 **quency**

226 A null hypothesis of no spatial pattern in allele frequencies can be generated
 227 by permuting the location of sampled populations among each other. Most
 228 empirical systems are not panmictic, and so this null model is trivial in a sense.
 229 However, comparison with this null across scales and locations can reveal when
 230 systems shift from small-scale homogeneity (from local gene flow) to larger scale
 231 heterogeneity (from limited gene flow) (Keitt 2007).

232 2.1.3 Simulated neutral patterns across a continuous landscape

233 To demonstrate the wavelet transformation of allele frequencies, and wavelet
 234 genetic dissimilarity function, we applied these tools to several simulated scen-
 235 arios. First, we conducted forward landscape genetic simulations under neu-
 236 trality using the SLiM software (Haller and Messer 2019), building off published
 237 approaches (C J Battey, Peter L Ralph, and Kern 2020). We simulated out-
 238 crossing, iteroparous, hermaphroditic organisms, with modest lifespans (aver-
 239 age of ~ 4 time steps). Individual fecundity was Poisson distributed, mating
 240 probability (determining paternity) was determined based on a Gaussian kernel
 241 (truncated at three standard deviations), and dispersal distance from mother
 242 was also Gaussian (C. Battey, Peter L Ralph, and Kern 2020). Individuals
 243 became mature in the time step following their dispersal. These parameters
 244 roughly approximate a short lived perennial plant with gene flow via pollen
 245 movement and seed dispersal. Competition reduced survival and decayed with
 246 distance following a Gaussian (truncated at three standard deviations, C J Bat-
 247 tey, Peter L Ralph, and Kern 2020). Near landscape boundaries, survival was
 248 reduced to compensate for lower competition from beyond the landscape mar-
 249 gin (C J Battey, Peter L Ralph, and Kern 2020). Code is available at GitHub
 250 (<https://github.com/jesserlasky/WaveletSpatialGenetic>).

251 We began by characterizing a simple scenario across a continuous land-
 252 scape. We simulated a square two dimensional landscape measuring 25 units
 253 on each side. The standard deviation of mating and dispersal distance σ were
 254 both 0.2, yielding a combined standard deviation of gene flow distances of 0.24
 255 $[(3\sigma^2/2)^{1/2}]$. In this first simulation there was no selection. The population was
 256 allowed to evolve for 100,000 time steps before we randomly sampled 200 indi-
 257 viduals and 1,000 SNPs with a minor allele frequency of at least 0.05. The first
 258 two principal components (PCs) of these SNPs show smooth population struc-
 259 ture across the landscape, and that these two PCs predict the spatial location
 260 of each sample (Figure S2 in File S1).

261 To facilitate interpretation of wavelet transformed allele frequencies
 262 $(T^{wav} f_i)(a, b, s)$ we provide two example loci i with distinct spatial patterns
 263 (Figure 1). The first locus has the greatest variance in wavelet transformed
 264 allele frequencies among sampled loci at $s = 0.4$ (Figure 1A-C) while the second
 265 locus has the greatest variance at $s = 12.2$ (Figure 1D-F).

266 We then calculated wavelet dissimilarity $D_{a,b}^{wav}(s)$, aggregating the signals
 267 in $(T^{wav} f_i)(a, b, s)$ across loci i , for each sampled location at a range of spatial
 268 scales s . Here and below we use a set of scales increasing by a constant log dis-
 269 tance interval, as genetic distances are often linearly correlated to log geographic
 270 distances in two dimensions (F. Rousset 1997). The mean across sampled lo-
 271 cations for each scale was calculated and compared to the null distribution for
 272 that scale (Figure S2 in File S1). The null was generated by permuting locations
 273 of sampled individuals as described above, and observed mean of dissimilarity
 274 was considered significant if it was below the 2.5 percentile or above the 97.5
 275 percentile of dissimilarity from null permutations.

276 When comparing our simulated data to the null, we found that mean wavelet




Figure 1: **Two example SNPs (rows) with distinct spatial patterns.** Shading shows either allelic variation (untransformed, A, D) or variation in wavelet transformed allele frequencies ($T^{wav} f_i$)(a, b, s) (B,C,E,F). The first locus (A-C) has the greatest variance in wavelet transformed allele frequency among sampled loci at $s = 0.4$. The second locus (D-F) has the greatest variance in wavelet transformed allele frequency at $s = 12.2$. For the SNP in the top row, the variance among locations in ($T^{wav} f_i$)(a, b, s) for $s = 0.4$ is 0.56 (visualized as shading in B), while it is only 0.17 for the SNP in the bottom row (E). For the SNP in the bottom row, the variance among locations in ($T^{wav} f_i$)(a, b, s) for $s = 12.2$ is 44.46 (visualized as shading in F), while it is only 1.24 for the SNP in the top row (C).

277 genetic dissimilarity was significantly less than expected under the null model
 278 at scales $s \leq 0.93$, due to local homogenization by gene flow (standard deviation
 279 = 0.24). At scales $s \geq 1.24$, wavelet dissimilarity was significantly greater than
 280 expected, due to isolation by distance, with monotonically increasing wavelet
 281 genetic dissimilarity at greater scales (Figure S2 in File S1).

282 To demonstrate how the scale of gene flow influences the wavelet dissimilarity
 283 $D_{a,b}^{wav}(s)$, we also conducted identical simulations as described above but instead
 284 with standard deviations of mating and dispersal distances, σ , of 0.5, 1, 2, or 5,
 285 yielding combined standard deviations of gene flow distances of 0.61, 1.22, 2.45,
 286 and 6.12.

287 To verify that simulations were generating results consistent with theoretical
 288 expectations of continuous populations at equilibrium, we compared the sim-
 289 ulated gene flow parameters with estimations from the simulated data based
 290 on theory. The slope of genetic differentiation versus geographic distance in
 291 two dimensions is expected to be proportional to the inverse of Wright’s neigh-
 292 borhood size, $4\pi D\sigma^2$, where D is the effective population density and σ is the
 293 standard deviation of gene flow (Rousset 2000; X. Vekemans and O. J. Hardy
 294 2004; Wright 1943, 1946).

295 We estimated D using $N_e = (4N - 2)/(V + 2)$ where N is census population
 296 size and V is variance in lifetime reproductive output (Kimura and Crow 1963).
 297 We calculated V using the lifetime reproductive output of the individuals dying
 298 in the last 50 time steps. We then divided the estimated N_e by landscape
 299 area (assuming evenly distribution across the landscape) to get effective density
 300 D (X. Vekemans and O. J. Hardy 2004). We used three different genetic
 301 differentiation or kinship metrics (Loiselle et al. 1995; Ritland 1996; Rousset
 302 2000) combined with estimated D to estimate gene flow across a range of true
 303 gene flow parameters (using SPAGeDi v1.5 software, Olivier J. Hardy and Xavier
 304 Vekemans (2002)). We also compared individual pairwise estimates of genetic
 305 differentiation across distance with the theoretically expected slope. Simulations
 306 were run for 100,000 time steps with parameters as described above.

307 We found that the gene flow estimated using the slope of genetic versus
 308 geographic distance and D was closely matched by the simulation parameter
 309 value, especially for the Rousset (2000) genetic differentiation estimator (Figures
 310 S3 and S4 in File S1). This matching suggests these simulations corresponded
 311 well with theory for continuous populations at equilibrium, despite ignoring the
 312 effects of negative density dependence, uneven distribution of individuals, and
 313 boundary effects (Nick H. Barton, Depaulis, and Etheridge 2002).

314 With increasing scale of gene flow we see a flatter change in wavelet dis-
 315 similarity across spatial scales (Figure S5 in File S1). When gene flow is local,
 316 wavelet dissimilarity is low at small scales and high at large scales. At the large
 317 gene flow scale, the observed wavelet dissimilarity is indistinguishable from the
 318 panmictic null. We also ran the same analyses but using biased sampling along
 319 the landscape’s y-axis, so that 3/4 of samples were in the upper half of the land-
 320 scape. Even with this bias, the wavelet dissimilarities across scales and gene flow
 321 parameters were essentially unchanged (Figure S6 in File S1). To investigate
 322 sensitivity to landscape size, we also ran these same simulations with landscapes

323 four times as large (50x50) and found similar patterns of wavelet dissimilarity
324 across scales and simulated gene flows (Figure S7 in File S1).

325 **3 Results**

326 **3.0.1 Simulated long-term neutral patterns in a heterogeneous land-** 327 **scape**

328 To assess if our approach could identify localized and scale-specific patterns of
329 isolation by distance, we next simulated multiple scenarios where we expected
330 spatial heterogeneity. First, we simulated neutral evolution across a simulated
331 patchy landscape (generated from earlier work) (Jesse R. Lasky and Keitt 2013).
332 This landscape contained a substantial portion of unsuitable habitat where ar-
333 riving propagules perished. We used the same population parameters as previ-
334 ously and simulated 100,000 time steps to reach approximately stable relatedness
335 patterns. We then calculated wavelet dissimilarity using 1,000 random SNPs of
336 200 sampled individuals.

337 Additionally, we sought to compare wavelet dissimilarities to more familiar
338 metrics. To do so, we calculated euclidean genetic distance (in the space of
339 allele frequencies across the genome) and geographic distance between pairs
340 of samples, and did this for different subsets of samples and regions, so as
341 to compare localized patterns in wavelet dissimilarity to localized patterns in
342 pairwise distances.

343 In our landscape, wavelet dissimilarity showed localized and scale-specific
344 patterns of low and high dissimilarity (Figure 2). Notably, the same two islands
345 (top left and bottom right of landscape in Figure 2) have lower dissimilarity
346 than expected at small scales and are more dissimilar than expected at larger
347 scales. Stated another way, these islands have low diversity locally (e.g. within
348 populations), as can be seen by the slow increase in genetic distance with geo-
349 graphic distance locally (Figure 2D, compare to 2F). However, at larger scales
350 (e.g. comparing island to mainland) islands are more dissimilar, as seen by the
351 greater genetic distances at larger geographic distances (Figure 2E, compare to
352 2G; also see the first two principal components of SNPs, Figure S8 in File S1).
353 These results highlight the capacity of the method to contrast patterns across
354 scales requiring only dilation of the analyzing kernel.

355 **3.0.2 Simulated neutral patterns in a colonizing and range-** 356 **expanding species**

357 For a second scenario where we expected localized, scale-specific heterogeneity,
358 we simulated an invasion/range expansion. Beyond the importance of invasions
359 in applied biology, the changes in spatial genetic patterns over time are of general
360 interest (Castric and Bernatchez 2003; Le Corre et al. 1997; Slatkin 1991, 1993),
361 considering that all species ranges are dynamic and many "native" species still
362 bear clear evidence of expansion, e.g. following the last glacial maximum.




Figure 2: **Wavelet genetic dissimilarity identifies scale-specific, localized patterns in a heterogeneous landscape, with pairwise distance plots for comparison.** (A-C) Maps of simulated landscape where habitat is gray (in background) and unsuitable areas are white. Sampled individuals are circles. Colors represent sampling locations where wavelet genetic dissimilarity was significantly high (red) or low (blue), with s , the wavelet scale, shown at top of each panel as a horizontal line. At the smallest scales (A), samples have less dissimilarity than expected, especially in the island in the upper left of the landscape. This pattern can also be seen (D,F) when comparing pairwise geographic versus euclidean genetic distances for samples in the different regions of the landscape (dashed grey lines in A). At larger spatial scales (B-C), all locations have significantly greater dissimilarity than expected due to limited gene flow. However, the same islands show the greatest dissimilarity at large scales (lower panels), due to their high genetic difference from mainland samples at center. This pattern can also be seen in the pairwise genetic distances across larger geographic distances (E,G). (D-G) Loess smoothing curves are shown.

363 We simulated invasion across a square landscape of the same size as before,
 364 but beginning with identical individuals only in the middle at the bottom edge
 365 of the landscape (Figure 3). We sampled 200 individuals at time steps 100, 250,
 366 500, 1000, 1500, 2000, through the full populating of the landscape around 2500
 367 years and until the 3000th time step.

368 We characterized wavelet genetic dissimilarity and found substantial hetero-
 369 geneity across different regions and across time (e.g. for $s = 6.9$, dark versus
 370 light red in Figure 3A-C). This heterogeneity in genetic turnover can be seen by
 371 contrasting genotypes from different regions. Near the expansion front, there
 372 is relative homogeneity and low diversity locally in new populations, but with
 373 rapid turnover in genotypes separated by space, resulting in high wavelet dis-
 374 similarity at intermediate spatial scales (Figure 3D). In the range interior, there
 375 is greater local diversity and less turnover in genotype across space, i.e. a weaker
 376 isolation by distance (Figure 3E, see all SNP genetic distance plot Figure S9 in
 377 File S1). Supporting the role of founder effects and low diversity at expanding
 378 range margins in driving these patterns, we observed a decline in medium- and
 379 large-scale wavelet dissimilarity in later years (Figure 3G) after the landscape
 380 had been populated.

381 These patterns highlight how wavelet dissimilarity is capturing scale-specific
 382 turnover in genetic composition, rather than merely genetic distance at a given
 383 geographic distance. Comparing the two regions highlighted in Figure 3B, the
 384 genetic distances at a geographic distance of 6.9 are not strikingly different
 385 (Figure S9 in File S1). Rather what distinguishes these regions is their rate
 386 of genetic change in composition at this scale, as highlighted in Figure 3. The
 387 region of high wavelet dissimilarity at $s = 6.9$ (Figure 3B) transitions from
 388 homogeneity among nearby samples to high genetic distance at larger scales
 389 (Figure 3D, S9). By contrast the region of low wavelet dissimilarity at $s = 6.9$
 390 (Figure 3B) starts out with greater genetic distance among nearby samples with
 391 a modest increase in genetic distance at larger scales (Figure 3E, S9).

392 Overall, these simulations show the capacity of $D_{a,b}^{wav}(s)$, wavelet genetic
 393 dissimilarity, to capture localized, scale specific trends in genetic composition.
 394 Given the spatial heterogeneity in nature and the dynamics of populations and
 395 species ranges through time, there are likely many such patterns waiting to be
 396 described to shed light on patterns of gene flow and population history.

397 **3.1 Finding the loci of local adaptation**

398 **3.1.1 Using wavelet transforms to identify outliers of spatial pattern** 399 **in allele frequency**

400 We can also use our approach to transforming allele frequencies to identify par-
 401 ticular genetic loci involved in local adaptation, and the regions and spatial
 402 scales of turnover in their allele frequency. Our strategy is (as before) to first
 403 calculate $(T^{wav} f_i)(a, b, s)$, the wavelet transform, for each locus i at each sam-
 404 pling point a, b for a set of chosen spatial scales $s \in S$.

405 Because of different ages and histories of drift, mutations will vary in their




Figure 3: **Wavelet genetic dissimilarity reveals dynamic spatial patterns during an invasion across a homogeneous landscape.** Left column of panels (A-C) shows a map of the landscape through time, with 200 sampled individuals at each time step and the wavelet dissimilarity at $s = 6.9$ at their location. Darker red indicates greater wavelet dissimilarity. In the second time step, 1000, two regions are highlighted in dashed boxes (B), one with higher dissimilarity at $s = 6.9$ (D) and one with lower dissimilarity at this scale (E). (D-E) show pairwise geographic distance versus distance in the first PC of SNPs for samples from these regions. (F) shows the loadings of each sample on the first PC of SNPs. (D-E) highlight the greater increase in PC1 distance with geographic distance at this scale (vertical dashed lines) in (D), compared to the smaller increase in PC1 distance at this scale in (E). In particular, the region highlighted in (D) is homogeneous at short distances but very distinct at distances at the highlighted scale $s = 6.9$, indicating the major genetic turnover at this scale and location. (G) Mean wavelet dissimilarity across the landscape changes over time, highlighting the dynamic spatial population genetic patterns across invasions. Loess smoothing curves are shown in (E-F).

406 global allele frequency and thus global variance. To facilitate comparisons
 407 among loci for relative evidence of selection, we can normalize spatial patterns in
 408 allele frequency by total variation across locations, as is done when calculating
 409 F_{ST} .

410 Here we divide the wavelet transforms of allele frequency by the standard
 411 deviation of global allele frequency variation for each locus i , $sd(f_i)$. This nor-
 412 malization is greatest when minor allele frequency is 0.5 for a biallelic locus, and
 413 yields a scaled wavelet transformed allele frequency: $(T^{wav} f_i)(a, b, s)/sd(f_i)$, for
 414 a given location and scale.

415 We then calculate the variance across sampling locations of
 416 $(T^{wav} f_i)(a, b, s)/sd(f_i)$ and refer to this quantity as the "scale-specific
 417 genetic variance." This scaled-specific variance is akin to F_{ST} in being a
 418 measure of spatial variation in allele frequency normalized to total variation
 419 (which is determined by mean allele frequency). High scale-specific variance
 420 for a given locus indicates high variation at that scale relative to the total
 421 variation and mean allele frequency. We then used a χ^2 null distribution across
 422 all genomic loci to calculate parametric p-values (Cavalli-Sforza 1966; Lewontin
 423 and Krakauer 1973) and used the approach of Whitlock and Lotterhos (2015) to
 424 fit the degrees of freedom of the distribution of scale-specific genetic variances
 425 (see Supplemental Methods). Applying this approach to a range of simulated
 426 scenarios as well as an empirical dataset (described below), we see that the χ^2
 427 distribution with a maximum likelihood fit to determine degrees of freedom
 428 provides a reasonably close fit to the distribution of scale-specific genetic
 429 variance among SNPs (Figures S10-S13 in File S1).

430 3.1.2 Simulated local adaptation

431 First, we present some specific individual simulations for illustration, and then
 432 a larger set with more variation in underlying parameters. We simulated a
 433 species with the same life history parameters as in simulations above, with
 434 the addition of spatially varying viability selection on a quantitative trait. We
 435 imposed two geometries of spatially varying selection, one a linear gradient
 436 and the other a square patch of different habitat selecting for a different trait
 437 value. As with the neutral simulations, simulations with selection began with
 438 organisms distributed across the landscape, with an ancestral trait value of
 439 zero. In these simulations, 1% of mutations influenced the quantitative trait
 440 with additive effects and with effect size normally distributed with a standard
 441 deviation of 5. For the linear gradient, the optimal trait value was 0.5 at one
 442 extreme and -0.5 at the other extreme, on a 25x25 square landscape. Selection
 443 was imposed using a Gaussian fitness function to proportionally reduce survival
 444 probability, with standard deviation σ_k . In this first simulation, $\sigma_k = 0.5$.
 445 Carrying capacity was roughly 5 individuals per square unit area, and simulated
 446 populations usually stabilized close to this density. Full details of simulation,
 447 including complete code, can be found in supplemental materials and on GitHub
 448 (<https://github.com/jesserlasky/WaveletSpatialGenetic>).

449 In the first simulation along a linear gradient after 2,000 time steps there

450 were 2 selected loci with minor allele frequency (MAF) at least 0.1, with a
 451 genetic variance in the trait of 3.7. (the scale of mating and propagule dispersal
 452 were each $\sigma = 1.1$) The two loci under stronger selection were clearly identified
 453 by the scale-specific genetic variance $var((T^{wav} f_i)(a, b, s)/sd(f_i))$ at the larger
 454 spatial scales (Figure 4). When there is a linear selective gradient across the
 455 entire landscape, the largest spatial scale is the one most strongly differentiating
 456 environments and the strongest scale-specific genetic variance was at the largest
 457 scale (Figure 4). However, power may not be greatest at these largest scales,
 458 because population structure also is greatest at these largest scales. Instead,
 459 power was greatest at intermediate scales, as seen by the lowest p-values being
 460 detected at these intermediate scales (Figure 4). At these scales there is greater
 461 gene flow but still some degree of changing selection that may maximize power
 462 to detect selection.

463 We next simulated change in selection in a discrete habitat patch, which
 464 may more closely correspond to the setting where researchers would find useful
 465 a flexible approach to finding spatial patterns in allele frequency, especially
 466 if the patches of distinct environment are not known by researchers. In our
 467 simulation there was a large central patch, 10x10, that selected for distinct trait
 468 values (trait optimum = 0.5) compared to the outer parts of the landscape (trait
 469 optimum = -0.5). Selection was initially weakly stabilizing ($\sigma_k = 3$ around the
 470 optimum of zero for the first 500 years to accumulate diversity, and then the
 471 patch selective differences were imposed with stronger selection, $\sigma_k = 0.08$. The
 472 scales of mating and propagule dispersal were each $\sigma = 2$. Carrying capacity
 473 was roughly 50 individuals per square unit area.

474 In this simulation we present results after 3000 time steps, where there was
 475 a single common QTL under selection, giving a genetic variance in trait of 0.42
 476 (Figure 5). We found several spurious large scale peaks in scale-specific genetic
 477 variance (Figure 5A), but when using the χ^2 test on these statistics we clearly
 478 identified the single QTL under selection, with lowest p-values for intermediate
 479 scales (Figure 5B).

480 We calculated the scale-specific genetic variance across a denser spectrum
 481 of scales s for the causal SNP, to determine at what scale variance was great-
 482 est. We found the maximum scale-specific genetic variance for the causal SNP
 483 was at 5.02, approximately half the length of a patch edge (Figure 5C). For
 484 illustration, we also calculated F_{ST} (Goudet 2005; Weir and Cockerham 1984)
 485 for several naively discretized subpopulation scenarios for a simple illustration
 486 of how results are sensitive to discretization (Figure 5D-F). We also implement
 487 our test on these two simulated landscapes but with biased sampling and found
 488 our ability to detect causal loci was robust (Figure S14 in File S1).

489 3.1.3 Evaluating the scale-specific genetic variance test

490 As an initial assessment of the general appropriateness of the scale-specific ge-
 491 netic variance test we proposed above, we conducted additional simulations on
 492 two types of landscapes with varying life history parameters. These simulations
 493 were not meant to be an exhaustive evaluation of the performance of this new




Figure 4: **Scale-specific genetic variance test applied to simulations with a linear selective gradient.** (top panels) Genome-wide variation in scale-specific genetic variance, $\text{var}((T^{wav} f_i)(a, b, s)/sd(f_i))$, for five different scales s and upper-tail p-values for χ^2 test using fitted values of d.f. Each point represents a SNP at a specific scale. Loci under selection are indicated with vertical lines along with the absolute value of the derived allele's effect on the trait and MAF. At bottom are shown maps of the two selected loci as well as their spectra of scale-specific genetic variance. At upper right the mean scale-specific genetic variance across all genomic loci is shown for each scale s . The scale of mating and propagule dispersal were each $\sigma = 1.1$. Gaussian viability selection was imposed with $\sigma_k = 0.5$. Carrying capacity was approximately 5 individuals per square unit area.




Figure 5: **Simulations of local adaptation to a single discrete patch of different habitat.** (A) Genome-wide variation in scale-specific genetic variance $var((T^{wav} f_i)(a, b, s)/sd(f_i))$ and (B) χ^2 p-values for six different scales s , for a discrete habitat difference after 3000 simulated years. Each point in the left panels represents a SNP, and wavelet statistics (A-B) at specific scales. The selected SNP is indicated with a vertical line along with the absolute value of a derived allele's effect on the trait and MAF. (C) A map of the landscape with individuals' genotypes at the causal SNP indicated with color, in addition to the spectrum of scale-specific genetic variance at this SNP, showing a peak at approximately half the patch width (vertical line at 5). (D-E) Implementation of F_{ST} using arbitrary boundaries for populations. This approach can easily miss causal loci (C,E) if the delineated population boundaries do not match habitat boundaries. (A) At upper right the mean scale-specific genetic variance across all loci is shown for each scale s . The scale of mating and propagule dispersal were each $\sigma = 2$. Gaussian viability selection was imposed with $\sigma_k = 0.08$.

494 test; we leave a more extensive evaluation for future studies.

495 Here, we again used the discrete habitat patch landscape and the linear
496 gradient landscape but with a wider range of parameter variation. We tested
497 a range of mating and dispersal (σ) scales including 0.25, 0.5, 1, and 2, and a
498 range of stabilizing selection (σ_k) values including 0.125, 0.25, 0.5, and 1. Three
499 simulations were conducted for each combination of parameter settings and each
500 ran for 10,000 years.

501 Because PCAdapt is one of the few methods for identification of spatial
502 pattern in allele frequency that does not require subpopulation discretization
503 and in theory could detect patterns at multiple scales, we also implemented
504 this method. We used the PCA of the scaled genotype matrix, thinned for
505 LD but including causal SNPs, to extract the z-scores and p-values of each
506 SNP with a cutoff of $p = 0.05$. We used a scree plot showing the percentage
507 of variance explained in decreasing order to identify the optimal number of
508 principal components following Cattell's rule (Duforet-Frebourg et al. 2016).

509 Calculating false and true positive rates for PCAdapt was straightforward,
510 but for the scale specific genetic variance test there are several tests (one at
511 each scale) for each SNP. To conservatively represent inference across these
512 multiple tests, we considered SNPs a significant result if one of the tested scales
513 was significant. Because the individual scale tests are slightly conservative,
514 and continuous wavelet transforms are correlated across scales (and hence not
515 completely independent tests), we expected the resulting false positive rates
516 would not be unduly high.

517 Overall the scale-specific genetic variance test showed good false positive
518 rates. Across simulations, the proportion of SNPs with χ^2 upper-tail $p < 0.05$
519 at one scale was usually close to but sometimes slightly more than 0.05 (Figure
520 6). By contrast, under scenarios of low gene flow and strong stabilizing selection,
521 nominal false positive rates were high for PCAdapt, often > 0.15 .

522 Power to detect SNPs (proportion of selected SNPs with $p < 0.05$) under
523 selection was generally high (true positive rate near 1) but sometimes low, de-
524 pending on the strength of selection (σ_k) and mating and dispersal scales (σ)
525 (Figure 6). When gene flow was high and selection was weak, power was low
526 for both the scale-specific genetic variance test and PCAdapt. This also corre-
527 sponds to the scenario when local adaptation is weakest (Kirkpatrick and N. H.
528 Barton 1997). In addition to considering power simply based on p for each SNP,
529 we also considered power using the top p -value rank among selected SNPs under
530 each simulation, based on the reasoning that researchers may want to follow up
531 on top ranked outlier SNPs first before any lower ranked SNPs. This approach
532 showed similar results, with high power for both the scale-specific genetic vari-
533 ance test and PCAdapt except when gene flow was high and selection weak. In
534 general, the two methods showed comparable power across different scenarios
535 (Figure 6), with some indication that the scale-specific genetic variance test had
536 higher power under high gene flow and PCAdapt slightly higher power under
537 lower gene flow. By plotting individual SNPs we can see that for the upper end
538 of gene flow scenarios ($\sigma = 1$ or 2), the scale-specific genetic variance test more
539 consistently identified selected SNPs at the top compared to PCAdapt. For the

540 low gene flow scenarios, PCAdapt more consistently identified large effect vari-
 541 ants, while the scale-specific genetic variance test more consistently identified
 542 the smaller effect variants (see results for linear gradient in Figure S15 in File
 543 S1). Overall, the similarities in true and false positive rates between methods
 544 suggest that our wavelet approach is effective compared to other related tools,
 545 while our test also offers the ability to explicitly consider variation in spatial
 546 scale.

547 **3.2 Testing for spatial pattern in quantitative trait loci** 548 **(QTL)**

549 When testing for spatially-varying selection on a quantitative trait, one approach
 550 is ask whether QTL identified from association or linkage mapping studies show
 551 greater allele frequency differences among populations than expected (Berg and
 552 G. Coop 2014; Price et al. 2018). Here we implement such an approach to
 553 compare wavelet transformed allele frequencies for QTL L to a set of randomly
 554 selected loci of the same number and distribution.

555 For this test we calculate the mean of scale-specific genetic variance for all
 556 QTL with MAF at least 0.05 among sampled individuals. We then permute
 557 the identity of causal QTL across the genome and recalculate the mean scale-
 558 specific genetic variance, and repeat this process 1000 times to generate a null
 559 distribution of mean scale-specific genetic variance of QTL for each scale s .

560 We illustrate this test here briefly using a simulation of adaptation to a
 561 square patch of habitat in the middle of a landscape, with the two gene flow
 562 parameters $\sigma = 0.5$, the strength of selection $\sigma_K = 0.5$, carrying capacity
 563 ~ 5 individuals per square unit area. After 1000 generations we sampled 300
 564 individuals, from which there were 13 QTL for the trait under selection with
 565 MAF at least 0.05. We then calculated the mean scale-specific genetic variance
 566 for these QTL across scales s and compared to the null permutations of randomly
 567 selected 13 SNPs from the genome.

568 We found significantly higher mean scale-specific genetic variance for the
 569 QTL than the null expectation at all 6 scales tested. Although the scale-specific
 570 genetic variance was greatest at the largest scales for the QTL, these scales did
 571 not show as great a distinction when comparing to the null. The greatest mean
 572 wavelet variance of QTL relative to null came at the intermediate scales of 3-5,
 573 which was approximately 1/3-1/2 the width of the habitat patch (Figure S16 in
 574 File S1).

575 **3.3 Application to an empirical system**

576 **3.3.1 Genome-wide wavelet dissimilarity**

577 We applied our approach to an empirical dataset of diverse, broadly distributed
 578 genotypes with whole genome resequencing data: 908 genotypes from 680 nat-
 579 ural populations of the model plant, *Arabidopsis thaliana* (Brassicaceae). We




Figure 6: Comparing the scale-specific genetic variance test with PCAadapt in simulations of adaptation to single discrete patch of different habitat. (A) True positive rates (nominal $p < 0.05$) for each combination of simulation parameters, the scales of mating and dispersal σ and the standard deviation of the Gaussian stabilizing selection function σ_k . (B) An alternate view of statistical power based on the median rank of the top selected SNP among all SNPs. (C) False positive rates (nominal $p < 0.05$). (D) Comparing power between the two statistical approaches for the different simulation runs. Density of points is shown in the blue scale so as to indicate where many simulations had the same result. The line indicates a 1:1 relationship. (E-F) Individual selected SNPs in simulations, showing their nominal p values and ranks among all SNPs, colored based on σ in the simulation. The x-axis represents the proportion of total phenotypic variation among sampled individuals that was explained by the given SNP (R^2 from a linear model).

580 used a published Arabidopsis dataset (Alonso-Blanco et al. 2016), only includ-
581 ing Eurasian populations and excluding highly distinct "relicts" and also likely
582 contaminant accessions (Pisupati et al. 2017). For locations with more than one
583 accession genotyped we calculated allele frequency. We used a total of 129536
584 SNPs filtered for minor allele frequency ($MAF > 0.05$) and LD (Zheng et al.
585 2012).

586 We first calculated the genome-wide wavelet dissimilarity, $D_{a,b}^{wav}(s)$, across a
587 series of increasing scales s at even intervals in log distance units from ~ 50 m
588 to approximately half the distance separating the farthest samples, ~ 3000 km.

589 We observed increasing mean genome-wide wavelet dissimilarity at larger
590 scales (Figure 7), a pattern indicative of isolation by distance, on average, across
591 the landscape. Arabidopsis showed significantly low dissimilarity at scales less
592 than ~ 5 km, likely due to the homogenizing effect of gene flow. However, we
593 found significantly high dissimilarity at scales greater than ~ 7 km. This scale
594 of significantly high dissimilarity may be a relatively short distance, consid-
595 ering that Arabidopsis is largely self pollinating and lacks clear seed dispersal
596 mechanisms (though seeds of some genotypes form mucus in water that increases
597 buoyancy) (Saez-Aguayo et al. 2014). At scales greater than ~ 120 km we found
598 an increase in the slope relating scale s and dissimilarity, perhaps signifying a
599 scale at which local adaptation begins to emerge.

600 The locations of scale-specific dissimilarity among Arabidopsis populations
601 revealed several interesting patterns. Even by the ~ 30 km scale, there were
602 three notable regions of significantly high dissimilarity: northern Spain and
603 extreme southern and northern Sweden (Figure 7). The high dissimilarity at this
604 scale in northern Spain corresponds to the most mountainous regions of Iberia,
605 suggesting that limitations to gene flow across this rugged landscape have led
606 to especially strong isolation among populations at short distances. In northern
607 Sweden, Long et al. (2013) previously found a particularly steep increase in
608 isolation-by-distance. Alonso-Blanco et al. (2016) found that genetic distance
609 was greatest among accessions from Southern Sweden at scales from $\sim 20 - 250$
610 compared to regions farther south. At larger, among-region scales, dissimilarity
611 was significantly high across the range, with Iberia and northern Sweden again
612 being most dissimilar at ~ 234 km and surpassed by central Asia at ~ 1834 km
613 as being most dissimilar. Iberia and northern Sweden contain many accessions
614 distantly related to other accessions, likely due to isolation during glaciation
615 and subsequent demographic histories (Alonso-Blanco et al. 2016). This scale
616 in Asia separates populations in Siberia from those further south in the Tian
617 Shan and Himalayas, indicating substantial divergence potentially due to limited
618 gene flow across the heterogeneous landscape. By contrast, populations in the
619 UK and the Balkan peninsula had low dissimilarity across a range of scales,
620 possibly due to reduced diversity and a more recent history of spread in these
621 regions.




Figure 7: **Genome-wide wavelet dissimilarity, $D_{a,b}^{wav}(s)$, for *Arabidopsis* genotypes.** (A) The global mean dissimilarity across scales compared to the null expectation (gray ribbon) and (B) the dissimilarity across scales centered on each sampled genotype, with several regions highlighted (vertical lines indicate scales shown in panels C-F). (C-F) Selected scales highlight the changes in dissimilarity across locations, with each circle indicating a genotyped sample/population. Red indicates significantly greater wavelet dissimilarity than expected, blue significantly less than expected. For the map panels, the intensity of color shading indicates the relative variation (for a given scale) in $D_{a,b}^{wav}(s)$ among significant locations.

622 3.3.2 Identifying putative locally-adapted loci

623 For this analysis, we used the same genotypes as in the prior section but not
 624 filtered for LD, leaving 1,642,040 SNPs with $MAF > 0.1$ (Alonso-Blanco et al.
 625 2016).

626 The scale-specific genetic variance test identified putative locally adapted
 627 loci (Figure S17 in File S1). The distribution of scale-specific genetic variance
 628 among SNPs was reasonably matched to the theoretical χ^2 distribution (Figure
 629 S13 in File S1). Among notable loci, at the ~ 59 km scale, the #2 QTL and #3
 630 SNP is in the coding region of METACASPASE 4 (MC4), a gene that controls
 631 biotic and abiotic stress-induced programmed cell death (Hander et al. 2019;
 632 Shen, Liu, and Li 2019). To speculate, if MC4 were involved in coevolution
 633 with microbial pathogens we might expect rapid allele frequency dynamics and
 634 thus a pattern of high variation among even nearby populations.

635 The #1 SNP for the ~ 282 km scale was in the coding sequence of the DOG1
 636 gene (Figure 8, Figure S17 in File S1). This SNP, Chr. 5, 18,590,741 was also
 637 strongly associated with flowering time (see next section) and germination and
 638 tags known functional polymorphisms at this gene that are likely locally adap-
 639 tive (Martínez-Berdeja et al. 2020). The spatial pattern of variation at this
 640 locus (Figure 8) is complicated, highlighting the benefit of the flexible wavelet
 641 approach. By contrast, imposing a grid on this landscape, or using national
 642 political boundaries to calculate F_{ST} could easily miss the signal as did Horton
 643 et al. (2012). The climate-allele frequency associations for DOG1 are also com-
 644 plicated and non-monotonic (**gamba'genomics'2023** ; Martínez-Berdeja et al.
 645 2020), making it challenging for genotype-environment association approaches
 646 (Jesse R Lasky, Emily B Josephs, and Morris 2023).

647 At the ~ 1359 km scale, the #1 SNP (and also the lowest p-value SNP among
 648 all scales, Figure 8, Figure S17 in File S1) was on chromosome 5 at 26,247,515 bp,
 649 555 bp upstream from AT5G65660, a hydroxyproline-rich glycoprotein family
 650 protein. These are cell wall glycoproteins with important roles in development
 651 and growth (Johnson et al. 2017) some of which have a role in abiotic stress
 652 response (Tseng et al. 2013).

653 3.3.3 Testing for local adaptation in quantitative trait loci (QTL)

654 We tested for non-random scale-specific genetic variance of QTL for Arabidopsis
 655 flowering time, a trait that is likely involved in local adaptation (Ågren et al.
 656 2017). We used previously published data on flowering time: days to flower at
 657 10°C measured on 1003 genotypes and days to flower at 16°C measured on 970
 658 resequenced genotypes (Alonso-Blanco et al. 2016). We then performed mixed-
 659 model genome wide association studies (GWAS) in GEMMA (v 0.98.3) (Zhou
 660 and Stephens 2012) with 2,048,993 M SNPs filtered for minor allele frequency
 661 ($MAF > 0.05$), while controlling for genome-wide similarity among ecotypes.

662 We found that top flowering time GWAS SNPs showed significantly elevated
 663 scale-specific genetic variance at several intermediate spatial scales tested. For
 664 flowering time at both 10° and 16°C , scale-specific genetic variance was signifi-




Figure 8: Allelic variation (colors) for SNPs that were top outliers for scale-specific genetic variance test at different scales. On maps at left, the scale for which a SNP was an outlier is indicated by a bar above each map. The right panels show the spatial spectra for each SNP, i.e. the scale-specific genetic variance across a range of scales. Dashed lines indicate the scale for which a SNP was an outlier.

665 cantly elevated for the top 1,000 SNPs at the 282, 619, and 1359 km scales, but
666 not always at the largest or smallest scales (Figure 9). In particular the scale-
667 specific genetic variances were greatest for the ~ 282 km scale where the mean
668 scale specific genetic variance for 16°C QTL was 15.2 standard deviations above
669 the null mean, and the ~ 619 km scale, where the mean scale specific genetic
670 variance for 10°C QTL was 13.5 standard deviations above the null mean. For
671 QTL from both temperature experiments, results were nearly equivalent if we
672 instead used the top 100 SNPs.

673 4 Discussion

674 Geneticists have long developed theory for spatial patterns in allele frequency
675 (Haldane 1948; Malécot 1948; Wright 1943). Empiricist have sought to use these
676 patterns make inference about underlying processes of demography, gene flow,
677 and selection (Lewontin and Krakauer 1973; McRae et al. 2008; Rousset 2000).
678 While statistical approaches have been developed to characterize geographic
679 patterns, few are flexible enough to incorporate patterns at a range of scales that
680 are also localized in space. Because wavelet transforms have these properties,
681 we think they may be useful tools for geneticists. Here we demonstrated several
682 applications of wavelet transforms to capture patterns in whole genome variation
683 and at particular loci, under a range of neutral and non-neutral scenarios.

684 Some important existing approaches are based on discretization of spatially-
685 distributed samples into spatial bins, i.e. putative populations (Bishop, Cham-
686 bers, and I. J. Wang 2023; Petkova, Novembre, and Stephens 2016; Weir and
687 Cockerham 1984). However, without prior knowledge of selective gradients, pat-
688 terns of gene flow, or relevant barriers, it is often unclear how to delineate these
689 populations. For example, we can see how the specific discretization can hinder
690 our ability to find locally-adapted loci in our simulations (Figure 5) and in em-
691 pirical studies of *Arabidopsis* in the case of the phenology gene *DOG1* that was
692 missed in previous F_{ST} scans (Alonso-Blanco et al. 2016; Horton et al. 2012).

693 Our goal in this paper was to provide a new perspective on spatial popula-
694 tion genetics using the population-agnostic, and spatially smooth approach of
695 wavelet transforms. We showed how these transforms characterize scale-specific
696 and localized population structure across landscapes (Figures 2, 3, 7). We also
697 showed how wavelet transforms can capture scale-specific evidence of selection
698 on individual genetic loci (Figures 4, 5, 6, 8) and on groups of quantitative trait
699 loci (Figure 9). Our simulations and empirical examples showed substantial
700 heterogeneity in the scale and stationarity of spatial patterns. For example,
701 the wavelet genetic dissimilarity allowed us to identify regions near a front of
702 range expansion with steeper isolation by distance at particular scales due to
703 drift (Figure 3). Additionally, we identified loci underlying local adaptation
704 and showed an example where the evidence for this adaptation was specific to
705 intermediate spatial scales (Figure 5). While existing approaches to character-
706 izing population structure or local adaptation have some ability to characterize
707 scale specific patterns, e.g. those based on ordinations of geography (Wagner,




Figure 9: **Testing for selection on *Arabidopsis* flowering time QTL.** We compared scale-specific genetic variance, $var((T^{wav} f_i)(a, b, s)/sd(f_i))$, of QTL with random SNPs, for five different scales s , for flowering time measured at 10°C and 16°C. The first two columns show the observed mean of the top 1,000 flowering time SNPs with a vertical line and a z-score. The histograms show null distributions of scale-specific genetic variance based on permutations of an equal number of markers with an equal distribution as the flowering time QTL. At right the scale-specific genetic variance is shown for random SNPs and for the flowering time QTL (gray lines), across scales, with the mean indicated by a black line.

708 Chávez-Pesqueira, and Forester 2017) or SNPs (Emily B. Josephs et al. 2019),
709 and some can capture localized patterns (e.g Petkova, Novembre, and Stephens
710 2016), there are few examples of approaches that merge both abilities (Wagner,
711 Chávez-Pesqueira, and Forester 2017).

712 Like many methods in population genetics that rely on inference from ob-
713 servational data, we view our approaches as exploratory and hypothesis gen-
714 erating. Heterogeneous patterns of genome-wide wavelet dissimilarity suggest
715 demographic hypotheses, some of which can be tested with detailed ecological
716 and genetic study (e.g. Keeley et al. 2017). For genome-scans for loci involved
717 in local adaptation, the p-values resulting from multiple tested scales are compa-
718 rable and so we recommend starting with the loci having the lowest p-value, and
719 using these to develop hypotheses for functional follow up experiments (Jesse R
720 Lasky, Emily B Josephs, and Morris 2023).

721 The test for spatial pattern in individual loci we developed owes greatly to
722 previous work from Lewontin and Krakauer (1973) who initially developed χ^2
723 tests applied to the distribution of F_{ST} values, and from Whitlock and Lotterhos
724 (2015)'s approach of inferring the degrees of freedom of the χ^2 distribution
725 using maximum likelihood and F_{ST} across loci. The χ^2 distribution underlies
726 a number of related genetic applied across loci (François et al. 2016). However,
727 we note that this test may be slightly conservative in some situations (Figure 6).
728 Nevertheless, we believe there were important signs in our work that this χ^2 -
729 based scale-specific genetic variance test was valuable. In particular, we found
730 in our simulation of adaptation to a habitat patch that the scale-specific genetic
731 variance was greatest at large spatial scales but at neutral sites, which obscured
732 spatial pattern at the causal locus (Figure 5). When applying the χ^2 test, we
733 were able to clearly map the causal locus while spurious loci with high scale-
734 specific genetic variance fell away because spatial patterns at those loci still fit
735 within the null distribution.

736 Relatedly, we found in other simulations and our empirical examples that the
737 strongest evidence for local adaptation was often not at the largest spatial scales
738 (Figure 9), even when the selective gradient was linear across the landscape (i.e.
739 the largest scale, Figure 4). This enhanced power at scales sometimes smaller
740 than the true selective gradients may be due to the limited power to resolve
741 true adaptive clines at large scales from the genome-wide signal of isolation by
742 distance at these scales. At intermediate scales, there may be a better balance
743 of sufficient environmental variation to generate spatial pattern versus higher
744 relatedness between locations due to gene flow.

745 We note that there remain several limitations to our approach proposed
746 here. First, the ability of wavelet transforms to capture patterns depends on
747 the correspondence between the wavelet form (shape) and the form of the em-
748 pirical patterns we seek to enhance, and there may be better functional forms to
749 filter spatial patterns in allele frequency. Generally speaking, a more compact
750 smoothing kernel with minimum weight in the tails will be better at reveal-
751 ing abrupt spatial transitions, but at the necessary cost of less precise deter-
752 mination of scale (Heisenberg 1927). Smoothing kernels such as the tricube

753 ($k_x \simeq [1 - x^3]^3$) have been shown to optimize certain trade-offs in this space
 754 and could be used to construct a difference-of-kernels wavelet. However, the
 755 overall influence of kernel shape tends to be much less than the influence of ker-
 756 nel bandwidth in our experience. Second, we have not yet implemented localized
 757 tests for selection (i.e. specific to certain locations) as we did with genome-wide
 758 dissimilarity. A challenge applying this test at individual loci is that there is a
 759 very large number of resulting tests from combinations of loci, locations, and
 760 scales. Therefore we have not fully exploited the localized information we derive
 761 from the wavelet transforms.

762 There are number of interesting future directions for research on wavelet
 763 characterization of spatial pattern in evolutionary biology. First, we could apply
 764 the wavelet transforms to genetic variation in quantitative traits measured in
 765 common gardens, to develop tests for selection on traits akin to the $Q_{ST} - F_{ST}$
 766 test (Emily B. Josephs et al. 2019; Whitlock and Guillaume 2009). Second, we
 767 could follow the example of Al-Asadi et al. (2019) and apply our measures of
 768 genetic dissimilarity to haplotypes of different size to estimate relative variation
 769 in the age of population structure. Third, we should test the performance of
 770 our tools under a wider range of demographic and selective scenarios to get a
 771 more nuanced picture of their strengths and weaknesses. Fourth, null models for
 772 wavelet dissimilarity could be constructed using knowledge of gene flow processes
 773 (instead of random permutation) to identify locations and scales with specific
 774 deviations from null patterns of gene flow.

775 4.1 Conclusion

776 Population genetics (like most fields) has a long history of arbitrary discretiza-
 777 tion for the purposes of mathematical, computational, and conceptual conve-
 778 nience. However, the real world often exists without clear boundaries between
 779 populations and where processes act simultaneously at multiple scales. We be-
 780 lieve that wavelet transforms are one of a range of tools that can move population
 781 genetics into a richer but still useful characterization of the natural world.

782 4.2 Data availability

783 Code used to generate the simulations and analyses shown here are freely avail-
 784 able at <https://github.com/jesserlasky/WaveletSpatialGenetic/>.

785 5 Acknowledgments

786 We thank Emily Josephs and Benjamin Peter and two anonymous reviewers for
 787 helpful comments. We thank Joanna Rifkin and JGI for help finding bugs in
 788 code.

789 6 Funding

790 This work was supported by NIH award R35GM138300 to JRL.

791 References

- 792 Ågren, Jon et al. (Mar. 2017). “Adaptive divergence in flowering time among
793 natural populations of *Arabidopsis thaliana*: Estimates of selection and QTL
794 mapping”. In: *Evolution* 71.3, pp. 550–564. ISSN: 0014-3820. DOI: 10.1111/
795 evo.13126. URL: <https://doi.org/10.1111/evo.13126> (visited on
796 08/28/2023).
- 797 Alonso-Blanco, Carlos et al. (2016). “1,135 Genomes Reveal the Global Pattern
798 of Polymorphism in *Arabidopsis thaliana*”. In: *Cell* 166, pp. 481–491. ISSN:
799 0092-8674. DOI: 10.1016/j.cell.2016.05.063.
- 800 Al-Asadi, Hussein et al. (Jan. 2019). “Estimating recent migration and
801 population-size surfaces”. en. In: *PLOS Genetics* 15.1. Publisher: Public Li-
802 brary of Science, e1007908. ISSN: 1553-7404. DOI: 10.1371/journal.pgen.
803 1007908. URL: [https://journals.plos.org/plosgenetics/article?id=
804 10.1371/journal.pgen.1007908](https://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1007908) (visited on 02/24/2022).
- 805 Ayton, Peter and Ilan Fischer (Dec. 2004). “The hot hand fallacy and the gam-
806 bler’s fallacy: Two faces of subjective randomness?” en. In: *Memory & Cog-
807 nition* 32.8, pp. 1369–1378. ISSN: 1532-5946. DOI: 10.3758/BF03206327.
808 URL: <https://doi.org/10.3758/BF03206327> (visited on 01/18/2024).
- 809 Barton, Nick H., Frantz Depaulis, and Alison M. Etheridge (Feb. 2002). “Neu-
810 tral Evolution in Spatially Continuous Populations”. In: *Theoretical Popu-
811 lation Biology* 61.1, pp. 31–48. ISSN: 0040-5809. DOI: 10.1006/tpbi.2001.
812 1557. URL: [https://www.sciencedirect.com/science/article/pii/
813 S0040580901915576](https://www.sciencedirect.com/science/article/pii/S0040580901915576) (visited on 01/21/2024).
- 814 Battey, C J, Peter L Ralph, and Andrew D Kern (May 2020). “Space is the
815 Place: Effects of Continuous Spatial Structure on Analysis of Population
816 Genetic Data”. In: *Genetics* 215.1, pp. 193–214. ISSN: 1943-2631. DOI: 10.
817 1534/genetics.120.303143. URL: [https://doi.org/10.1534/genetics.
818 120.303143](https://doi.org/10.1534/genetics.120.303143) (visited on 01/28/2024).
- 819 — (June 2020). “Predicting geographic location from genetic variation with
820 deep neural networks”. In: *eLife* 9. Ed. by Patricia J Wittkopp et al. Pub-
821 lisher: eLife Sciences Publications, Ltd, e54507. ISSN: 2050-084X. DOI: 10.
822 7554/eLife.54507. URL: <https://doi.org/10.7554/eLife.54507> (vis-
823 ited on 01/25/2022).
- 824 Berg, Jeremy J. and Graham Coop (Aug. 2014). “A Population Genetic Signal
825 of Polygenic Adaptation”. In: *PLoS Genet* 10.8, e1004412. DOI: 10.1371/
826 journal.pgen.1004412. URL: [http://dx.doi.org/10.1371%2Fjournal.
827 pgen.1004412](http://dx.doi.org/10.1371%2Fjournal.pgen.1004412).
- 828 Bhatia, Gaurav et al. (Sept. 2013). “Estimating and interpreting FST: The im-
829 pact of rare variants”. en. In: *Genome Research* 23.9. Company: Cold Spring
830 Harbor Laboratory Press Distributor: Cold Spring Harbor Laboratory Press

831 Institution: Cold Spring Harbor Laboratory Press Label: Cold Spring Har-
832 bor Laboratory Press Publisher: Cold Spring Harbor Lab, pp. 1514–1521.
833 ISSN: 1088-9051, 1549-5469. DOI: 10.1101/gr.154831.113. URL: <http://genome.cshlp.org/content/23/9/1514> (visited on 08/06/2020).

834 Bishop, Anusha P., E. Anne Chambers, and Ian J. Wang (2023). “Gen-
835 erating continuous maps of genetic diversity using moving win-
836 dows”. en. In: *Methods in Ecology and Evolution* 14.5. eprint:
837 [https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1111/2041-](https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1111/2041-210X.14090)
838 [210X.14090](https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1111/2041-210X.14090), pp. 1175–1181. ISSN: 2041-210X. DOI: 10.1111/2041-
839 [210X.14090](https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1111/2041-210X.14090). URL: [https://onlinelibrary.wiley.com/doi/abs/10.](https://onlinelibrary.wiley.com/doi/abs/10.1111/2041-210X.14090)
840 [1111/2041-210X.14090](https://onlinelibrary.wiley.com/doi/abs/10.1111/2041-210X.14090) (visited on 01/29/2024).

841 Blakemore, S.-J. et al. (Nov. 2003). “The detection of intentional contingencies
842 in simple animations in patients with delusions of persecution”. en. In: *Psy-*
843 *chological Medicine* 33.8. Publisher: Cambridge University Press, pp. 1433–
844 1441. ISSN: 1469-8978, 0033-2917. DOI: 10.1017/S0033291703008341.
845 URL: [https://www.cambridge.org/core/journals/psychological-](https://www.cambridge.org/core/journals/psychological-medicine/article/detection-of-intentional-contingencies-in-simple-animations-in-patients-with-delusions-of-persecution/EB6864B67E97778980DBB47A91279AB6)
846 [medicine/article/detection-of-intentional-contingencies-in-](https://www.cambridge.org/core/journals/psychological-medicine/article/detection-of-intentional-contingencies-in-simple-animations-in-patients-with-delusions-of-persecution/EB6864B67E97778980DBB47A91279AB6)
847 [simple-animations-in-patients-with-delusions-of-persecution/](https://www.cambridge.org/core/journals/psychological-medicine/article/detection-of-intentional-contingencies-in-simple-animations-in-patients-with-delusions-of-persecution/EB6864B67E97778980DBB47A91279AB6)
848 [EB6864B67E97778980DBB47A91279AB6](https://www.cambridge.org/core/journals/psychological-medicine/article/detection-of-intentional-contingencies-in-simple-animations-in-patients-with-delusions-of-persecution/EB6864B67E97778980DBB47A91279AB6) (visited on 01/18/2024).

849 Bradburd, Gideon S., Graham M. Coop, and Peter L. Ralph (Sept. 2018).
850 “Inferring Continuous and Discrete Population Genetic Structure Across
851 Space”. eng. In: *Genetics* 210.1, pp. 33–52. ISSN: 1943-2631. DOI: 10.1534/
852 [genetics.118.301333](https://doi.org/10.1534/genetics.118.301333).

853 Bradburd, Gideon S. and Peter L. Ralph (2019). “Spatial Population Genetics:
854 It’s About Time”. In: *Annual Review of Ecology, Evolution, and System-*
855 *atics* 50.1. eprint: <https://doi.org/10.1146/annurev-ecolsys-110316-022659>,
856 pp. 427–449. DOI: 10.1146/annurev-ecolsys-110316-022659. URL:
857 <https://doi.org/10.1146/annurev-ecolsys-110316-022659> (visited on
858 01/25/2024).

859 Castric, Vincent and Louis Bernatchez (Mar. 2003). “The rise and fall of iso-
860 lation by distance in the anadromous brook charr (*Salvelinus fontinalis*
861 Mitchill).” In: *Genetics* 163.3, pp. 983–996. ISSN: 0016-6731. URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1462472/> (visited on
862 11/28/2022).

863 Cavalli-Sforza, L. L. (Mar. 1966). “Population structure and human evolution”.
864 In: *Proceedings of the Royal Society of London. Series B. Biological Sciences*
865 164.995. Publisher: Royal Society, pp. 362–379. DOI: 10.1098/rspb.1966.
866 0038. URL: [http://royalsocietypublishing.org/doi/10.1098/rspb.](http://royalsocietypublishing.org/doi/10.1098/rspb.1966.0038)
867 [1966.0038](http://royalsocietypublishing.org/doi/10.1098/rspb.1966.0038) (visited on 02/10/2022).

868 Daubechies, Ingrid (1992). *Ten lectures on wavelets*. SIAM. ISBN: 0-89871-274-2.

869 Duforet-Frebourg, Nicolas et al. (Apr. 2016). “Detecting Genomic Signatures
870 of Natural Selection with Principal Component Analysis: Application to the
871 1000 Genomes Data”. In: *Molecular Biology and Evolution* 33.4, pp. 1082–
872 1093. ISSN: 0737-4038. DOI: 10.1093/molbev/msv334. URL: [https://doi.](https://doi.org/10.1093/molbev/msv334)
873 [org/10.1093/molbev/msv334](https://doi.org/10.1093/molbev/msv334) (visited on 02/24/2022).

874
875

- 876 Excoffier, Laurent and Nicolas Ray (July 2008). “Surfing during population
877 expansions promotes genetic revolutions and structuration”. eng. In: *Trends*
878 *in Ecology & Evolution* 23.7, pp. 347–351. ISSN: 0169-5347. DOI: 10.1016/
879 *j.tree*.2008.04.004.
- 880 Fitzpatrick, Matthew C. and Stephen R. Keller (2015). “Ecological genomics
881 meets community-level modelling of biodiversity: mapping the genomic land-
882 scape of current and future environmental adaptation”. In: *Ecology Letters*
883 18.1, pp. 1–16. ISSN: 1461-0248. DOI: 10.1111/ele.12376. URL: <http://dx.doi.org/10.1111/ele.12376>.
- 884 François, Olivier et al. (2016). “Controlling false discoveries in genome
885 scans for selection”. en. In: *Molecular Ecology* 25.2. _eprint:
886 <https://onlinelibrary.wiley.com/doi/pdf/10.1111/mec.13513>, pp. 454–469.
887 ISSN: 1365-294X. DOI: 10.1111/mec.13513. URL: [http://onlinelibrary-](http://onlinelibrary-wiley-com/doi/abs/10.1111/mec.13513)
888 [wiley-com/doi/abs/10.1111/mec.13513](http://onlinelibrary-wiley-com/doi/abs/10.1111/mec.13513) (visited on 02/11/2022).
- 889 Frantz, A. C. et al. (2009). “Using spatial Bayesian methods to deter-
890 mine the genetic structure of a continuously distributed population:
891 clusters or isolation by distance?” en. In: *Journal of Applied Ecol-*
892 *ogy* 46.2. _eprint: [https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-](https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-2664.2008.01606.x)
893 [2664.2008.01606.x](https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-2664.2008.01606.x), pp. 493–505. ISSN: 1365-2664. DOI: 10.1111/j.1365-
894 [2664.2008.01606.x](https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-2664.2008.01606.x). URL: [https://onlinelibrary.wiley.com/doi/abs/](https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2664.2008.01606.x)
895 [10.1111/j.1365-2664.2008.01606.x](https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2664.2008.01606.x) (visited on 01/26/2024).
- 896 Fyfe, Sophie et al. (Nov. 2008). “Apophenia, theory of mind and schizotypy:
897 Perceiving meaning and intentionality in randomness”. In: *Cortex*. Special
898 Issue on “Neuropsychology of Paranormal Experiences and Beliefs” 44.10,
899 pp. 1316–1325. ISSN: 0010-9452. DOI: 10.1016/j.cortex.2007.07.
900 009. URL: [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S0010945208001391)
901 [S0010945208001391](https://www.sciencedirect.com/science/article/pii/S0010945208001391) (visited on 01/18/2024).
- 902 Gamba, Diana et al. (Aug. 2023). *The genomics and physiology of abiotic stres-*
903 *sors associated with global elevation gradients in Arabidopsis thaliana*. en.
904 Pages: 2022.03.22.485410 Section: New Results. DOI: 10.1101/2022.03.22.
905 485410. URL: [https://www.biorxiv.org/content/10.1101/2022.03.22.](https://www.biorxiv.org/content/10.1101/2022.03.22.485410v3)
906 [485410v3](https://www.biorxiv.org/content/10.1101/2022.03.22.485410v3) (visited on 12/07/2023).
- 907 Gautier, Mathieu (Dec. 2015). “Genome-Wide Scan for Adaptive Divergence
908 and Association with Population-Specific Covariates”. In: *Genetics* 201.4,
909 pp. 1555–1579. ISSN: 0016-6731. DOI: 10.1534/genetics.115.181453. URL:
910 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4676524/> (visited on
911 07/07/2021).
- 912 Goudet, Jérôme (2005). “hierfstat, a package for r to compute and test
913 hierarchical F-statistics”. en. In: *Molecular Ecology Notes* 5.1. _eprint:
914 <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1471-8286.2004.00828.x>,
915 pp. 184–186. ISSN: 1471-8286. DOI: 10.1111/j.1471-8286.2004.00828.x.
916 URL: [http://onlinelibrary-wiley-com/doi/abs/10.1111/j.1471-](http://onlinelibrary-wiley-com/doi/abs/10.1111/j.1471-8286.2004.00828.x)
917 [8286.2004.00828.x](http://onlinelibrary-wiley-com/doi/abs/10.1111/j.1471-8286.2004.00828.x) (visited on 02/10/2022).
- 918 Groh, Jeffrey and Graham Coop (May 2023). “The temporal and genomic scale
919 of selection following hybridization”. In: *bioRxiv*, p. 2023.05.25.542345. DOI:
920

- 921 10.1101/2023.05.25.542345. URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10276902/> (visited on 01/26/2024).
- 922
- 923 Haldane, J. B. S. (Jan. 1948). “The theory of a cline”. eng. In: *Journal of*
924 *Genetics* 48.3, pp. 277–284. ISSN: 0022-1333. DOI: 10.1007/BF02986626.
- 925 Haller, Benjamin C. and Philipp W. Messer (2019). “SLiM 3: Forward Genetic
926 Simulations Beyond the Wright–Fisher Model”. en. In: *Molecular Biology*
927 *and Evolution* 36.3, pp. 632–637. DOI: 10.1093/molbev/msy228. URL: <http://academic.oup.com/mbe/advance-article/doi/10.1093/molbev/msy228/5229931> (visited on 01/19/2019).
- 928
- 929
- 930 Hancock, Zachary B., Rachel H. Toczydlowski, and Gideon S. Bradburd (Mar.
931 2023). “A spatial approach to jointly estimate Wright’s neighborhood size
932 and long-term effective population size”. In: *bioRxiv*, p. 2023.03.10.532094.
933 DOI: 10.1101/2023.03.10.532094. URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10029013/> (visited on 01/22/2024).
- 934
- 935 Hander, Tim et al. (Mar. 2019). “Damage on plants activates Ca²⁺-dependent
936 metacaspases for release of immunomodulatory peptides”. eng. In: *Science*
937 *(New York, N.Y.)* 363.6433, eaar7486. ISSN: 1095-9203. DOI: 10.1126/
938 *science.aar7486*.
- 939 Hardy, Olivier J. and Xavier Vekemans (2002). “spagedi: a versatile com-
940 puter program to analyse spatial genetic structure at the individual
941 or population levels”. en. In: *Molecular Ecology Notes* 2.4. eprint:
942 <https://onlinelibrary.wiley.com/doi/pdf/10.1046/j.1471-8286.2002.00305.x>,
943 pp. 618–620. ISSN: 1471-8286. DOI: 10.1046/j.1471-8286.2002.00305.x.
944 URL: <https://onlinelibrary.wiley.com/doi/abs/10.1046/j.1471-8286.2002.00305.x> (visited on 01/21/2024).
- 945
- 946 Heisenberg, W. (Mar. 1927). “Über den anschaulichen Inhalt der quanten-
947 theoretischen Kinematik und Mechanik”. de. In: *Zeitschrift für Physik*
948 43.3. Company: Springer Distributor: Springer Institution: Springer Label:
949 Springer Number: 3 Publisher: Springer-Verlag, pp. 172–198. DOI: 10.1007/
950 BF01397280. URL: <https://link.springer.com/article/10.1007/BF01397280> (visited on 03/21/2022).
- 951
- 952 Horton, Matthew W et al. (2012). “Genome-wide patterns of genetic variation
953 in worldwide *Arabidopsis thaliana* accessions from the RegMap panel”. In:
954 *Nat Genet* 44, pp. 212–216. ISSN: 1546-1718. DOI: 10.1038/ng.1042. URL:
955 <http://dx.doi.org/10.1038/ng.1042>.
- 956 Johnson, Kim L. et al. (June 2017). “Insights into the Evolution of
957 Hydroxyproline-Rich Glycoproteins from 1000 Plant Transcriptomes”. In:
958 *Plant Physiology* 174.2, pp. 904–921. ISSN: 0032-0889. DOI: 10.1104/pp.17.00295. URL: <https://doi.org/10.1104/pp.17.00295> (visited on
959 08/24/2023).
- 960
- 961 Josephs, Emily B. et al. (Mar. 2019). “Detecting Adaptive Differentiation in
962 Structured Populations with Genomic Data and Common Gardens”. en. In:
963 *Genetics* 211.3. Publisher: Genetics Section: Investigations, pp. 989–1004.
964 ISSN: 0016-6731, 1943-2631. DOI: 10.1534/genetics.118.301786. URL:
965 <https://www.genetics.org/content/211/3/989> (visited on 11/18/2020).

- 966 Kawecki, Tadeusz J. and Dieter Ebert (2004). “Conceptual issues in local adap-
 967 tation”. In: *Ecology Letters* 7.12, pp. 1225–1241. ISSN: 1461-0248. DOI: 10 .
 968 1111/j.1461-0248.2004.00684.x. URL: [http://dx.doi.org/10.1111/j.
 969 1461-0248.2004.00684.x](http://dx.doi.org/10.1111/j.1461-0248.2004.00684.x).
- 970 Keeley, Annika T. H. et al. (May 2017). “Habitat suitability is a poor proxy for
 971 landscape connectivity during dispersal and mating movements”. In: *Land-
 972 scape and Urban Planning* 161, pp. 90–102. ISSN: 0169-2046. DOI: 10.1016/
 973 j.landurbplan.2017.01.007. URL: [https://www.sciencedirect.com/
 974 science/article/pii/S0169204617300154](https://www.sciencedirect.com/science/article/pii/S0169204617300154) (visited on 02/01/2024).
- 975 Keitt, Timothy H. (2007). “On the quantification of local variation in biodiver-
 976 sity scaling using wavelets”. In: *Scaling Biodiversity*. Publisher: Cambridge
 977 University Press, pp. 168–80.
- 978 Kimura, Motoo and James F. Crow (1963). “The Measure-
 979 ment of Effective Population Number”. en. In: *Evolution* 17.3.
 980 _eprint: [https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1558-
 981 5646.1963.tb03281.x](https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1558-5646.1963.tb03281.x), pp. 279–288. ISSN: 1558-5646. DOI: 10.1111/j.1558-
 982 5646.1963.tb03281.x. URL: [https://onlinelibrary.wiley.com/doi/
 983 abs/10.1111/j.1558-5646.1963.tb03281.x](https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1558-5646.1963.tb03281.x) (visited on 01/20/2024).
- 984 Kimura, Motoo and George H. Weiss (Apr. 1964). “The Stepping Stone Model
 985 of Population Structure and the Decrease of Genetic Correlation with Dis-
 986 tance”. In: *Genetics* 49.4, pp. 561–576. ISSN: 0016-6731. URL: [https://www.
 987 ncbi.nlm.nih.gov/pmc/articles/PMC1210594/](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1210594/) (visited on 01/22/2024).
- 988 Kirkpatrick, Mark and N. H. Barton (July 1997). “Evolution of a Species’
 989 Range”. In: *The American Naturalist* 150.1. ArticleType: research-article
 990 / Full publication date: July 1997 / Copyright © 1997 The University of
 991 Chicago Press, pp. 1–23. ISSN: 00030147. URL: [http://www.jstor.org/
 992 stable/10.1086/286054](http://www.jstor.org/stable/10.1086/286054).
- 993 Lasky, Jesse R, Emily B Josephs, and Geoffrey P Morris (Jan. 2023). “Geno-
 994 type–environment associations to reveal the molecular basis of environmental
 995 adaptation”. In: *The Plant Cell* 35.1, pp. 125–138. ISSN: 1040-4651. DOI: 10 .
 996 1093/plcell/koac267. URL: [https://doi.org/10.1093/plcell/koac267
 997 \(visited on 02/26/2023\)](https://doi.org/10.1093/plcell/koac267).
- 998 Lasky, Jesse R., DAVID L. Des Marais, et al. (2012). “Characterizing genomic
 999 variation of *Arabidopsis thaliana*: the roles of geography and climate”. In:
 1000 *Molecular Ecology* 21.22, pp. 5512–5529. ISSN: 1365-294X. DOI: 10.1111/j .
 1001 1365-294X.2012.05709.x. URL: [http://dx.doi.org/10.1111/j.1365-
 1002 294X.2012.05709.x](http://dx.doi.org/10.1111/j.1365-294X.2012.05709.x).
- 1003 Lasky, Jesse R. and Timothy H. Keitt (Nov. 2013). “Reserve Size and Fragmenta-
 1004 tion Alter Community Assembly, Diversity, and Dynamics.” In: *The Amer-
 1005 ican Naturalist* 182.5, E142–E160. ISSN: 0003-0147. DOI: 10.1086/673205.
 1006 URL: <http://dx.doi.org/10.1086/673205> (visited on 02/22/2017).
- 1007 Le Corre, Valérie et al. (Apr. 1997). “Colonization with long-distance seed
 1008 dispersal and genetic structure of maternally inherited genes in forest
 1009 trees: a simulation study”. en. In: *Genetics Research* 69.2. Publisher:
 1010 Cambridge University Press, pp. 117–125. ISSN: 1469-5073, 0016-6723.
 1011 DOI: 10 . 1017 / S0016672397002668. URL: <https://www.cambridge>.

- 1012 org / core / journals / genetics - research / article / colonization -
 1013 with - longdistance - seed - dispersal - and - genetic - structure - of -
 1014 maternally - inherited - genes - in - forest - trees - a - simulation - study /
 1015 FFEDD8A5B195F1C01C99261FE49B6432 (visited on 11/28/2022).
- 1016 Lewontin, R. C. and J. Krakauer (May 1973). "Distribution of gene frequency
 1017 as a test of the theory of the selective neutrality of polymorphisms". eng. In:
 1018 *Genetics* 74.1, pp. 175–195. ISSN: 0016-6731.
- 1019 — (June 1975). "Testing the Heterogeneity of F Values". In: *Genetics* 80.2,
 1020 pp. 397–398. ISSN: 0016-6731. URL: [https://www.ncbi.nlm.nih.gov/pmc/
 1021 articles/PMC1213337/](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1213337/) (visited on 02/10/2022).
- 1022 Loiseau, Bette A. et al. (1995). "Spatial Genetic Structure of a Tropical Under-
 1023 story Shrub, *Psychotria officinalis* (Rubiaceae)". In: *American Journal
 1024 of Botany* 82.11. Publisher: Botanical Society of America, pp. 1420–1425.
 1025 ISSN: 0002-9122. DOI: 10.2307/2445869. URL: [https://www.jstor.org/
 1026 stable/2445869](https://www.jstor.org/stable/2445869) (visited on 01/21/2024).
- 1027 Long, Quan et al. (Aug. 2013). "Massive genomic variation and strong selection
 1028 in *Arabidopsis thaliana* lines from Sweden". en. In: *Nature Genetics*
 1029 45.8. Bandiera_abtest: a Cg.type: Nature Research Journals Number: 8 Primary_atype:
 1030 Research Publisher: Nature Publishing Group Subject_term: Population genetics
 1031 Subject_term_id: population-genetics, pp. 884–890. ISSN: 1546-1718. DOI: 10.1038/ng.2678. URL: [http://www-nature-com/
 1032 articles/ng.2678](http://www-nature-com/articles/ng.2678) (visited on 01/16/2022).
- 1033 Machado, Heather E et al. (June 2021). "Broad geographic sampling reveals
 1034 the shared basis and environmental correlates of seasonal adaptation in
 1035 *Drosophila*". In: *eLife* 10. Ed. by Magnus Nordborg, Patricia J Wittkopp,
 1036 and Magnus Nordborg. Publisher: eLife Sciences Publications, Ltd, e67577.
 1037 ISSN: 2050-084X. DOI: 10.7554/eLife.67577. URL: [https://doi.org/10.
 1038 7554/eLife.67577](https://doi.org/10.7554/eLife.67577) (visited on 01/24/2022).
- 1039 Malécot, G. (1948). *Les mathématiques de l'hérédité*. fre. OCLC: 832598694.
 1040 Paris: Masson & Cie.
- 1041 Manel, Stéphanie et al. (Apr. 2003). "Landscape genetics: combining landscape
 1042 ecology and population genetics". en. In: *Trends in Ecology & Evolution*
 1043 18.4, pp. 189–197. ISSN: 0169-5347. DOI: 10.1016/S0169-5347(03)00008-
 1044 9. URL: [https://www.sciencedirect.com/science/article/pii/
 1045 S0169534703000089](https://www.sciencedirect.com/science/article/pii/S0169534703000089) (visited on 10/16/2022).
- 1046 Martínez-Berdeja, Alejandra et al. (Feb. 2020). "Functional variants of DOG1
 1047 control seed chilling responses and variation in seasonal life-history strategies
 1048 in *Arabidopsis thaliana*". en. In: *Proceedings of the National Academy of
 1049 Sciences* 117.5. Publisher: National Academy of Sciences Section: Biological
 1050 Sciences, pp. 2526–2534. ISSN: 0027-8424, 1091-6490. DOI: 10.1073/pnas.
 1051 1912451117. URL: <https://www.pnas.org/content/117/5/2526> (visited
 1052 on 08/20/2021).
- 1053 McRae, Brad H. et al. (2008). "Using Circuit Theory to Model Connectivity
 1054 in Ecology, Evolution, and Conservation". en. In: *Ecology* 89.10.
 1055 _eprint: [https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.1890/07-
 1056 1861.1](https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.1890/07-1861.1), pp. 2712–2724. ISSN: 1939-9170. DOI: 10.1890/07-1861.1. URL:

- 1058 <https://onlinelibrary.wiley.com/doi/abs/10.1890/07-1861.1>
1059 (visited on 08/27/2023).
- 1060 McVean, Gil (Oct. 2009). “A Genealogical Interpretation of Principal Components Analysis”. en. In: *PLOS Genetics* 5.10. Publisher: Public Library of
1061 Science, e1000686. ISSN: 1553-7404. DOI: 10.1371/journal.pgen.1000686.
1062 URL: [https://journals.plos.org/plosgenetics/article?id=10.1371/
1063 journal.pgen.1000686](https://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1000686) (visited on 10/14/2022).
- 1064 Muraki, S. (June 1995). “Multiscale volume representation by a DoG wavelet”.
1065 In: *IEEE Transactions on Visualization and Computer Graphics* 1.2. Confer-
1066 ence Name: IEEE Transactions on Visualization and Computer Graphics,
1067 pp. 109–116. ISSN: 1941-0506. DOI: 10.1109/2945.468408.
- 1068 Nagylaki, Thomas (Oct. 1978). “A diffusion model for geographically structured
1069 populations”. en. In: *Journal of Mathematical Biology* 6.4, pp. 375–382. ISSN:
1070 1432-1416. DOI: 10.1007/BF02463002. URL: [https://doi.org/10.1007/
1071 BF02463002](https://doi.org/10.1007/BF02463002) (visited on 01/22/2024).
- 1072 Nei, Masatoshi and Takeo Maruyama (June 1975). “Lewontin-Krakauer test for
1073 neutral genes”. In: *Genetics* 80.2, p. 395. ISSN: 0016-6731. URL: [https:
1074 //www.ncbi.nlm.nih.gov/pmc/articles/PMC1213335/](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1213335/) (visited on
1075 02/10/2022).
- 1076 Peter, Benjamin M, Desislava Petkova, and John Novembre (Apr. 2020). “Ge-
1077 netic Landscapes Reveal How Human Genetic Diversity Aligns with Geog-
1078 raphy”. In: *Molecular Biology and Evolution* 37.4, pp. 943–951. ISSN: 0737-
1079 4038. DOI: 10.1093/molbev/msz280. URL: [https://doi.org/10.1093/
1080 molbev/msz280](https://doi.org/10.1093/molbev/msz280) (visited on 01/09/2022).
- 1081 Peterman, William E. (2018). “ResistanceGA: An R package for the optimiza-
1082 tion of resistance surfaces using genetic algorithms”. en. In: *Methods in Ecol-
1083 ogy and Evolution* 9.6, pp. 1638–1647. ISSN: 2041-210X. DOI: 10.1111/2041-
1084 210X.12984. URL: [https://besjournals.onlinelibrary.wiley.com/
1085 doi/abs/10.1111/2041-210X.12984](https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1111/2041-210X.12984) (visited on 04/14/2019).
- 1086 Petkova, Desislava, John Novembre, and Matthew Stephens (Jan. 2016). “Visu-
1087 alizing spatial population structure with estimated effective migration sur-
1088 faces”. en. In: *Nature Genetics* 48.1. Bandiera_abtest: a Cg.type: Nature Re-
1089 search Journals Number: 1 Primary_atype: Research Publisher: Nature Pub-
1090 lishing Group Subject.term: Population genetics;Software Subject_term.id:
1091 population-genetics;software, pp. 94–100. ISSN: 1546-1718. DOI: 10.1038/
1092 ng.3464. URL: <http://www-nature-com/articles/ng.3464> (visited on
1093 09/10/2021).
- 1094 Pisupati, Rahul et al. (Dec. 2017). “Verification of Arabidopsis stock collections
1095 using SNPmatch, a tool for genotyping high-plexed samples”. In: *Scientific
1096 Data* 4, p. 170184. ISSN: 2052-4463. DOI: 10.1038/sdata.2017.184. URL:
1097 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5744633/> (visited on
1098 10/01/2018).
- 1099 Price, Nicholas et al. (May 2018). “Combining population genomics and fitness
1100 QTLs to identify the genetics of local adaptation in *Arabidopsis thaliana*”.
1101 en. In: *Proceedings of the National Academy of Sciences* 115.19, pp. 5028–
1102

- 1103 5033. ISSN: 0027-8424, 1091-6490. DOI: 10.1073/pnas.1719998115. URL:
 1104 <http://www.pnas.org/content/115/19/5028> (visited on 06/06/2018).
- 1105 Pugach, Irina et al. (2011). “Dating the age of admixture via wavelet trans-
 1106 form analysis of genome-wide data”. en. In: *Genome Biology* 12.2. Publisher:
 1107 BMC, R19. DOI: 10.1186/gb-2011-12-2-r19. URL: [https://www.ncbi.
 1108 nlm.nih.gov/pmc/articles/PMC3188801/](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3188801/) (visited on 01/26/2024).
- 1109 Ritland, Kermit (Apr. 1996). “Estimators for pairwise relatedness and indi-
 1110 vidual inbreeding coefficients”. en. In: *Genetics Research* 67.2. Publisher:
 1111 Cambridge University Press, pp. 175–185. ISSN: 1469-5073, 0016-6723. DOI:
 1112 10.1017/S0016672300033620. URL: [https://www.cambridge.org/
 1113 core/journals/genetics-research/article/estimators-for-
 1114 pairwise-relatedness-and-individual-inbreeding-coefficients/
 1115 9AE218BF6BF09CCCE18121AA63561CF7](https://www.cambridge.org/core/journals/genetics-research/article/estimators-for-pairwise-relatedness-and-individual-inbreeding-coefficients/9AE218BF6BF09CCCE18121AA63561CF7) (visited on 01/21/2024).
- 1116 Robertson, Alan (June 1975). “Remarks on the Lewontin-Krakauer test”. In:
 1117 *Genetics* 80.2, p. 396. ISSN: 1943-2631. DOI: 10.1093/genetics/80.2.
 1118 396. URL: <https://doi.org/10.1093/genetics/80.2.396> (visited on
 1119 02/10/2022).
- 1120 Rosenberg, Noah A. et al. (Dec. 2005). “Clines, Clusters, and the Effect of
 1121 Study Design on the Inference of Human Population Structure”. en. In:
 1122 *PLOS Genetics* 1.6. Publisher: Public Library of Science, e70. ISSN: 1553-
 1123 7404. DOI: 10.1371/journal.pgen.0010070. URL: [https://journals.
 1124 plos.org/plosgenetics/article?id=10.1371/journal.pgen.0010070](https://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.0010070)
 1125 (visited on 01/26/2024).
- 1126 Rousset (2000). “Genetic differentiation between individu-
 1127 als”. en. In: *Journal of Evolutionary Biology* 13.1. eprint:
 1128 <https://onlinelibrary.wiley.com/doi/pdf/10.1046/j.1420-9101.2000.00137.x>,
 1129 pp. 58–62. ISSN: 1420-9101. DOI: 10.1046/j.1420-9101.2000.00137.x.
 1130 URL: [https://onlinelibrary.wiley.com/doi/abs/10.1046/j.1420-
 1131 9101.2000.00137.x](https://onlinelibrary.wiley.com/doi/abs/10.1046/j.1420-9101.2000.00137.x) (visited on 01/21/2024).
- 1132 Rousset, François (Apr. 1997). “Genetic Differentiation and Estimation of Gene
 1133 Flow from F-Statistics Under Isolation by Distance”. en. In: *Genetics* 145.4.
 1134 Publisher: Oxford Academic, pp. 1219–1228. DOI: 10.1093/genetics/145.
 1135 4.1219. URL: <https://dx.doi.org/10.1093/genetics/145.4.1219>
 1136 (visited on 08/27/2023).
- 1137 Saez-Aguayo, Susana et al. (Mar. 2014). “Local Evolution of Seed Flotation
 1138 in Arabidopsis”. en. In: *PLOS Genetics* 10.3. Publisher: Public Library of
 1139 Science, e1004221. ISSN: 1553-7404. DOI: 10.1371/journal.pgen.1004221.
 1140 URL: [https://journals.plos.org/plosgenetics/article?id=10.1371/
 1141 journal.pgen.1004221](https://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1004221) (visited on 02/13/2022).
- 1142 Serre, David and Svante Pääbo (Sept. 2004). “Evidence for Gradients of Human
 1143 Genetic Diversity Within and Among Continents”. In: *Genome Research*
 1144 14.9, pp. 1679–1685. ISSN: 1088-9051. DOI: 10.1101/gr.2529604. URL:
 1145 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC515312/> (visited on
 1146 01/26/2024).
- 1147 Shen, Wenzhong, Jiuer Liu, and Jian-Feng Li (Nov. 2019). “Type-II Metacasp-
 1148 ases Mediate the Processing of Plant Elicitor Peptides in Arabidopsis”. In:

- 1149 *Molecular Plant* 12.11, pp. 1524–1533. ISSN: 1674-2052. DOI: 10.1016/j.
 1150 molp.2019.08.003. URL: [https://www.sciencedirect.com/science/
 1151 article/pii/S1674205219302643](https://www.sciencedirect.com/science/article/pii/S1674205219302643) (visited on 08/28/2023).
- 1152 Shirk, Andrew J. and Samuel A. Cushman (2014). “Spatially-explicit estima-
 1153 tion of Wright’s neighborhood size in continuous populations”. In: *Fron-
 1154 tiers in Ecology and Evolution* 2. ISSN: 2296-701X. URL: [https://www.
 1155 frontiersin.org/articles/10.3389/fevo.2014.00062](https://www.frontiersin.org/articles/10.3389/fevo.2014.00062) (visited on
 1156 01/22/2024).
- 1157 Slatkin, Montgomery (Oct. 1991). “Inbreeding coefficients and coalescence
 1158 times”. en. In: *Genetical Research* 58.2, pp. 167–175. ISSN: 0016-6723, 1469-
 1159 5073. DOI: 10.1017/S0016672300029827. URL: [https://www.cambridge.
 1160 org/core/product/identifier/S0016672300029827/type/journal_
 1161 article](https://www.cambridge.org/core/product/identifier/S0016672300029827/type/journal_article) (visited on 11/28/2022).
- 1162 — (1993). “Isolation by Distance in Equilibrium and Non-Equilibrium Popu-
 1163 lations”. In: *Evolution* 47.1. Publisher: [Society for the Study of Evolution,
 1164 Wiley], pp. 264–279. ISSN: 0014-3820. DOI: 10.2307/2410134. URL: [http:
 1165 //www.jstor.org/stable/2410134](http://www.jstor.org/stable/2410134) (visited on 10/16/2022).
- 1166 Tseng, I-Chieh et al. (Sept. 2013). “Abscisic Acid- and Stress-Induced Highly
 1167 Proline-Rich Glycoproteins Regulate Root Growth in Rice”. In: *Plant Phys-
 1168 iology* 163.1, pp. 118–134. ISSN: 0032-0889. DOI: 10.1104/pp.113.217547.
 1169 URL: <https://doi.org/10.1104/pp.113.217547> (visited on 08/24/2023).
- 1170 Vekemans, X. and O. J. Hardy (2004). “New insights from fine-scale spatial
 1171 genetic structure analyses in plant populations”. en. In: *Molecular Ecol-
 1172 ogy* 13.4. _eprint: [https://onlinelibrary.wiley.com/doi/pdf/10.1046/j.1365-
 1173 294X.2004.02076.x](https://onlinelibrary.wiley.com/doi/pdf/10.1046/j.1365-294X.2004.02076.x), pp. 921–935. ISSN: 1365-294X. DOI: 10.1046/j.1365-
 1174 294X.2004.02076.x. URL: [https://onlinelibrary.wiley.com/doi/abs/
 1175 10.1046/j.1365-294X.2004.02076.x](https://onlinelibrary.wiley.com/doi/abs/10.1046/j.1365-294X.2004.02076.x) (visited on 01/20/2024).
- 1176 Wagner, Helene H., Mariana Chávez-Pesqueira, and Brenna R. Forester
 1177 (2017). “Spatial detection of outlier loci with Moran eigenvec-
 1178 tor maps”. en. In: *Molecular Ecology Resources* 17.6. _eprint:
 1179 <https://onlinelibrary.wiley.com/doi/pdf/10.1111/1755-0998.12653>,
 1180 pp. 1122–1135. ISSN: 1755-0998. DOI: 10.1111/1755-0998.12653.
 1181 URL: [https://onlinelibrary.wiley.com/doi/abs/10.1111/1755-
 1182 0998.12653](https://onlinelibrary.wiley.com/doi/abs/10.1111/1755-0998.12653) (visited on 01/23/2022).
- 1183 Wang, Ian J., Wesley K. Savage, and H. Bradley Shaffer (2009).
 1184 “Landscape genetics and least-cost path analysis reveal unex-
 1185 pected dispersal routes in the California tiger salamander (*Am-
 1186 bystoma californiense*)”. en. In: *Molecular Ecology* 18.7. _eprint:
 1187 <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-294X.2009.04122.x>,
 1188 pp. 1365–1374. ISSN: 1365-294X. DOI: 10.1111/j.1365-294X.2009.04122.
 1189 x. URL: [http://onlinelibrary-wiley-com/doi/abs/10.1111/j.1365-
 1190 294X.2009.04122.x](http://onlinelibrary-wiley-com/doi/abs/10.1111/j.1365-294X.2009.04122.x) (visited on 10/16/2022).
- 1191 Wang, Jianan et al. (Jan. 2020). “Genomic signatures of seed mass adap-
 1192 tation to global precipitation gradients in sorghum”. en. In: *Heredity*
 1193 124.1. Bandiera_abtest: a Cc.license_type: cc_by Cg.type: Nature Research
 1194 Journals Number: 1 Primary_atype: Research Publisher: Nature Publish-

- 1195 ing Group Subject_term: Agricultural genetics;Evolutionary biology Sub-
 1196 ject_term_id: agricultural-genetics;evolutionary-biology, pp. 108–121. ISSN:
 1197 1365-2540. DOI: 10.1038/s41437-019-0249-4. URL: [http://www-nature-](http://www-nature-com/articles/s41437-019-0249-4)
 1198 [com/articles/s41437-019-0249-4](http://www-nature-com/articles/s41437-019-0249-4) (visited on 01/05/2022).
- 1199 Weir, B. S. and C. Clark Cockerham (1984). “Estimating F-Statistics for the
 1200 Analysis of Population Structure”. In: *Evolution* 38.6. Publisher: [Society
 1201 for the Study of Evolution, Wiley], pp. 1358–1370. ISSN: 0014-3820. DOI:
 1202 10.2307/2408641. URL: <https://www.jstor.org/stable/2408641> (visited
 1203 on 08/06/2020).
- 1204 Whitlock, Michael C. and Frederic Guillaume (Nov. 2009). “Testing for Spa-
 1205 tially Divergent Selection: Comparing QST to FST”. In: *Genetics* 183.3,
 1206 pp. 1055–1063. ISSN: 0016-6731. DOI: 10.1534/genetics.108.099812. URL:
 1207 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2778959/> (visited on
 1208 08/07/2020).
- 1209 Whitlock, Michael C. and Katie E. Lotterhos (2015). “Reliable Detection of
 1210 Loci Responsible for Local Adaptation: Inference of a Null Model through
 1211 Trimming the Distribution of FST”. In: *The American Naturalist* 186.S1.
 1212 Publisher: [The University of Chicago Press, The American Society of Nat-
 1213 uralists], S24–S36. ISSN: 0003-0147. DOI: 10.1086/682949. URL: <https://www.jstor.org/stable/10.1086/682949> (visited on 08/30/2021).
- 1214 Wright, Sewall (Mar. 1931). “EVOLUTION IN MENDELIAN POPULA-
 1215 TIONS”. In: *Genetics* 16.2, pp. 97–159. ISSN: 1943-2631. DOI: 10.1093/
 1216 [genetics/16.2.97](https://doi.org/10.1093/genetics/16.2.97). URL: <https://doi.org/10.1093/genetics/16.2.97>
 1217 (visited on 01/22/2024).
- 1218 — (Mar. 1943). “Isolation by Distance”. In: *Genetics* 28.2, pp. 114–138. ISSN:
 1219 0016-6731. URL: [https://www.ncbi.nlm.nih.gov/pmc/articles/](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1209196/)
 1220 [PMC1209196/](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1209196/) (visited on 01/21/2024).
- 1221 — (Jan. 1946). “Isolation by Distance under Diverse Systems of Mating”. In:
 1222 *Genetics* 31.1, pp. 39–59. ISSN: 0016-6731. URL: [https://www.ncbi.nlm.](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1209315/)
 1223 [nih.gov/pmc/articles/PMC1209315/](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1209315/) (visited on 01/21/2024).
- 1224 — (Jan. 1949). “The genetical structure of populations”. In: *Annals of Eugenics*
 1225 15.1. Publisher: John Wiley & Sons, Ltd, pp. 323–354. ISSN: 2050-1420. DOI:
 1226 10.1111/j.1469-1809.1949.tb02451.x. URL: [http://onlinelibrary-](http://onlinelibrary-wiley-com/doi/abs/10.1111/j.1469-1809.1949.tb02451.x)
 1227 [wiley-com/doi/abs/10.1111/j.1469-1809.1949.tb02451.x](http://onlinelibrary-wiley-com/doi/abs/10.1111/j.1469-1809.1949.tb02451.x) (visited on
 1228 08/06/2020).
- 1229 Yang, Wen-Yun et al. (June 2012). “A model-based approach for analysis of
 1230 spatial structure in genetic data”. In: *Nat Genet* 44.6, pp. 725–731. ISSN:
 1231 1061-4036. DOI: 10.1038/ng.2285. URL: [http://dx.doi.org/10.1038/](http://dx.doi.org/10.1038/ng.2285)
 1232 [ng.2285](http://dx.doi.org/10.1038/ng.2285).
- 1233 Yeaman, Sam et al. (2016). “Convergent local adaptation to climate in distantly
 1234 related conifers”. In: *Science* 353.6306, pp. 1431–1433.
- 1235 Zheng, Xiuwen et al. (Dec. 2012). “A high-performance computing toolset for
 1236 relatedness and principal component analysis of SNP data”. en. In: *Bioinfor-*
 1237 [matics](https://academic.oup.com/bioinformatics/article/28/24/3326/245844) 28.24. Publisher: Oxford Academic, pp. 3326–3328. ISSN: 1367-4803.
 1238 DOI: 10.1093/bioinformatics/bts606. URL: [http://academic.oup.com/](http://academic.oup.com/bioinformatics/article/28/24/3326/245844)
 1239 [bioinformatics/article/28/24/3326/245844](http://academic.oup.com/bioinformatics/article/28/24/3326/245844) (visited on 04/13/2020).
- 1240

1241 Zhou, Xiang and Matthew Stephens (July 2012). “Genome-wide efficient mixed-
1242 model analysis for association studies”. In: *Nat Genet* 44.7, pp. 821–824.
1243 ISSN: 1061-4036. DOI: 10.1038/ng.2310. URL: [http://dx.doi.org/10.](http://dx.doi.org/10.1038/ng.2310)
1244 [1038/ng.2310](http://dx.doi.org/10.1038/ng.2310).